

GRAPH OF MAPS

by
Sóstenes Lins

A thesis presented to the University of Waterloo
in partial fulfillment of the
requirements for the degree of
Doctor of philosophy in Mathematics

Waterloo, Ontario, Canada, 1980

©Sóstenes Lins

I hereby declare that I am the sole author of this thesis.

I authorize the University of Waterloo to lend this thesis to other institutions or individuals for the purpose of scholarly research.

Signature:

I further authorize the University of Waterloo to reproduce this thesis by photocopying or by other means, in total or in part, at the request of other institutions or individuals for the purpose of scholarly research.

Signature:

The University of Waterloo requires the signatures of all persons using or photocopying this thesis. Please sign below, and give address and date.

A meus Pais

Abstract

This work studies certain aspects of graphs embedded on surfaces. Initially, a colored graph model for a map of a graph on a surface is developed. Then, a concept analogous to (and extending) planar graph is introduced in the same spirit as planar abstract duality, and is characterized topologically. An extension of the Gauss code problem treating together the cases in which the surface involved is the plane or the real projective plane is established. The problem of finding a minimum transversal of orientation-reversing circuits in graphs on arbitrary surfaces is proved to be NP-complete and is algorithmically solved for the special case where the surface is the real projective plane.

ACKNOWLEDGEMENTS

I am indebted to Bernardete, my wife, whose moral support was invaluable in all phases of this work; to Lauro, my son, whose birth during this period filled my life with joy and thus greatly contributed to the relief of the unavoidable tensions.

I am grateful to my supervisor, Professor Daniel H. Younger. He sparked my curiosity towards the subject of graphs on surfaces. Our discussions, his suggestions and his always firm, patient, and friendly criticism, greatly influenced not only the thesis, but my attitude towards mathematics in general. I learned from every aspect of our relationship.

I want to thank my colleagues and friends Arnaldo Mandel and Ephraim Korach for innumerable stimulating mathematical discussions. I also thank Professor Herb Shank: some of the key ideas in this work arose from his theory on left-right paths and from our personal interaction. To Professor U.S.R. Murty I express my appreciation for sharing with me the enthusiasm for the graph colored model. His feedback was important to me.

I thank UFPE and to CAPES, in Brazil, for financial support. The CIDA/COMBRA project also provided some support. Finally, I wish to thank Mrs. Sue Embro for her excellent typing of the thesis, as well as for her patience with the corrections.

Chapter 0

INTRODUCTION

This thesis is about graphs embedded in compact surfaces satisfying the property that the complement of the graph in the surface is (topologically) a collection of disjoint open discs, called faces. We summarize the latter restriction by saying that the graph is well-embedded in the surface. A pair (G, S) where graph G is well-embedded in surface S is called a topological map of a graph or, for short, a t -map.

In Tutte [25] an algebraic abstract model for the above object is introduced and called simply “map”. In Chapter 1 of our work we define a map as a finite, cubic, 3-edge-colored graph in which two of the three colors induce squares.

The main point of the first part of Chapter 1 is to show that the three objects are equivalent. Indeed, we present a natural bijection between our maps and the algebraic maps of Tutte, which we denote for short as a -maps. We present also a natural bijection between our maps and the t -maps. With these bijections, an intuitive geometrical interpretation of Tutte’s axioms is established.

The correspondence between the maps and the a -maps is based on the construction of the Cayley color graph of the action of three fixed point free involutions, which are slightly different from the three permutations which

constitute Tutte's a -maps.

The correspondence between the maps and t -maps is based on the dual of the barycentric division of the t -map, a very simple topological operation. Specifically it consists of the thickening of each edge, which is replaced by a bounding digon on the surface, followed by the expansion of each vertex, yielding a cubic graph in which the edges of the original are replaced by squares.

A suitable 3-coloration of the edges of the expanded graph can be used to reconstruct the t -map; this is the basic idea: to encode the surface and the cyclic sequences of edges around vertices using three colors.

Fundamental concepts of t -maps can be directly interpreted in the 3-edge-colored cubic graph. For instance, the Euler characteristic is defined as a signed sum of the numbers of the three types of 2-colored polygons. The orientability of the surface, as we show in Theorem 1.6, is equivalent to the bipartiteness of the cubic graph. Therefore, since it is simpler, we take the latter as our definition of orientability. We use these definitions to give a new proof of the known fact that the Euler characteristic of an orientable surface is even (Theorem 1.7). The interpretation of the topological concept of an orientation-reversing circuit (a circuit in the graph of a t -map which has, in the surface, a neighborhood homeomorphic to the Moebius strip) is the simpler concept of a circuit with an odd number of edges. This is important in the proof of Lemma 3.4, a central fact in Chapter 3.

Theorem 1.8 is a combinatorial counterpart, in the 3-colored model, of the topological fact that the union of a set of vertex-disjoint orientation-reversing circuits in a t -map is not the boundary of a subset of faces. This fact is used to establish the result of Chapter 4. It is also used at the end of Chapter 2 (Theorem 2.14) to derive an upper bound for the number of independent vertices in certain graphs.

In the final part of Chapter 1 we introduce the structure of maps which forms the basis of Chapter 2. Some topological consequences of this structure

are also considered.

The reentrant paths in a planar t -map which alternate using the rightmost and leftmost edges at each vertex have been studied by Shank [24], who names them left-right paths. They are also known as Petrie polygons; see Coxeter [3]. These objects have natural generalization for arbitrary surfaces, and we name them, in this context, zigzag paths.

In our model the abstract equivalence between the following three cyclic sequences arising from t -maps becomes apparent:

- (a) the cyclic sequence of edges around a vertex;
- (b) the cyclic sequence of the edges in a boundary of a face;
- (c) the cyclic sequence of the edges in a zigzag path.

The symmetry between (a) and (b) is widely known. It is given by the dual t -map which interchanges vertices and faces. We observe that the dual maintains the zigzags. We introduce the “phial” map, whose corresponding t -map interchanges (relative to the original t -map) the vertices and the zigzags, maintaining the faces. Also, in the same way, we introduce the “antimap”, which interchanges the faces and the zigzags, maintaining the vertices.

The operations of taking the dual and taking the phial generate, from a given map, a set of six maps in correspondence with the elements of the symmetric group on three elements. This structure is used (Theorem 1.9) to prove that every graph in which the edges are doubled can be embedded so as to form a self-dual t -map.

Chapter 2 introduces and investigates a concept analogous to planar duality.

A theorem by Whitney [27] states that a graph is planar if and only if it has an abstract dual. This dual can be taken to mean another graph with the same edges as the first graph such that a cycle in one of them corresponds to a coboundary in the other.

We introduce the following concept: a graph is called (cycle) rich if there are two other graphs with the same edges as the first graph and such that a cycle in one of the three graphs corresponds to the symmetric difference of two coboundaries, one for each of the other two graphs. Thus, we have an abstract correspondence between cycles and coboundaries of three graphs instead of two.

The main result of the first part of the chapter shows (Theorem 2.3) that triples of graphs sharing the above relation are induced naturally from certain classes of t -maps. If a rich graph arises in this way, then we say that the graph is map-rich. We do not know whether there are rich graphs which are not map-rich. If not, this would complete the analogy with Whitney's Theorem.

It follows, by a corollary of Proposition 2.6, that every planar graph is rich (in fact map-rich). Hence, the concept of rich graph is not only analogous to the concept of planar graph, but is in fact a generalization.

The central fact, which makes the theory symmetric in the three graphs, Theorem 2.5. It states that for arbitrary t -maps, the intersection of the vector space (over $GF(2)$) of the coboundaries of its graph and the vector space generated by the boundaries of its faces is contained in the vector space generated by the cycles induced by its zigzag paths. In each one of the six maps mentioned before, the above spaces occur differently permuted. The six maps induce the three graphs, each one occurring twice. The symmetry follows from properties.

The main results of the second part of the chapter are of map-rich graphs (Theorem 2.9). The first of these states that a graph is map-rich if it has a special kind of embedding (where the vertices are represented by closed curves and the edges by intersection of the curves) in a surface whose Euler characteristic and orientability are specified functions of the graph. The second characterization states that a graph is map-rich if and only if the special kind of embedding induces in some surface, a (usual) t -map which is 2-face-colorable and such that the cycle space of its (4-regular) graph is

generated by the boundaries of its faces and the curves which represent the vertices of the original graph.

In the third part of the chapter we use the second characterization mentioned above to prove that every complete graph is map-rich (Theorems 2.11 and 2.12). Moreover, from this characterization is evident an easy topological “recipe” to produce map-rich graphs from t -maps with arbitrary surface, in the same sense that Whitney’s theorem permits the presentation of examples of abstract duality simply by the drawing of planar graphs. The first characterization of map-rich graphs in Theorem 2.9 is used (Lemma 2.13) to prove that some graphs are not map-rich, for instance, the graph obtained from the Petersen graph by doubling all its edges.

Trying to get more insight in the membership decision problem for map-rich graphs we were led to study the Gauss code problem, which is the topic of our next chapter.

The Gauss code problem is the following: given a cyclic sequence of symbols in which each symbol occurs twice, find necessary and sufficient conditions to embed a closed curve in the plane with the restriction that the cyclic sequence of self-intersections of the curve reproduces the given cyclic sequence, when the symbols are identified with the self-intersection points.

A brief survey of this problem can be found in Grünbaum [10-pp.72-73]. Lovasz and Marx [17] present a solution in terms of a forbidden substructure. Rosenstiehl [19] presents a solution in terms of a graph called an interlace graph, which is easily computable from the given cyclic sequence. This solution includes a polynomial algorithm to test and realize the embedding (if it exists).

A generalized version of this problem can be considered: dropping the planar restriction, we ask for a surface of minimum connectivity for which there is an embedding of a curve satisfying the prescribed sequence of self-intersections.

The embedding of the curve induces a t -map with a 4-regular graph, since each crossing point is visited twice. Consider the special case of the generalized problem, where the restriction of 2-face-colorability of the induced t -map is imposed.

Note that this restricted form is yet a generalization of the original problem, since every 4-regular graph embedded in the plane is 2-face-colorable. The motivation for considering this special case is that an algorithmic solution to it would enable us to determine whether an arbitrary cubic graph is map-rich.

The main result of Chapter 3 is an algorithmic answer for the question of embeddability satisfying the 2-colorability restriction in the case of surfaces of connectivity at most one. Our approach treats together the cases of the plane and projective plane. Particularized to the plane, our conditions reproduce those of Rosenstiehl [19].

This result uses some algebraic properties of t -maps with one vertex, which we establish. These properties are based on the generalization for arbitrary surfaces, of some homomorphisms, which we call crossing functions, introduced for the plane by Shank [24] and also used by Rosenstiehl [19].

The t -maps induced by the restricted form of the generalized Gauss code problem are in 1-1 correspondence with the t -maps with one vertex. A t -map with one vertex induces naturally a crossing function. The central fact, from which we specialize the conditions for the plane and the projective plane, is the equality (Theorem 3.3(a)) between the dimension of the image of the composition of the crossing functions induced by a map and its antimap (whose associated t -maps have one vertex), and the connectivity of the surface associated with the phial map.

Chapter 4 presents an algorithmic solution for the problem of finding a minimum transversal of orientation reversing circuits in graphs embedded in the real projective plane.

Our interest in this problem arises from the fact that for graphs embedded in arbitrary surfaces, it includes the problem of finding a maximum coboundary in an arbitrary graph (Proposition 4.2). Since the latter problem is known to be *NP*-complete, (Garey and Johnson [6]) so is the problem that we treat, if considered for graphs in arbitrary surfaces.

Certain subsets of faces of a t -map which has an eulerian graph, called reducers, are recognized. By means of a suitable operation which eliminates a reducer, a t -map is transformed to a “smaller” one, without change in the cardinality of a minimum transversal of orientation reversing circuits. For the case of graphs embedded in the projective plane, the non-existence of a reducer implies that the t -map has a simple form, looking like a system of lines in the real projective plane. From this fact we deduce the following minimax relation (Theorem 4.5): for an eulerian graph G well-embedded in the real projective plane, the cardinality of a minimum transversal of orientation-reversing circuits in G is equal to the cardinality of a maximum disjoint collection of orientation-reversing circuits in G .

An important fact, which permits the use of the reducers, is that for projective t -maps a minimal transversal of orientation-reversing circuits is an orientation-reversing circuit in the dual t -map. We generalize this relation by showing in Theorem 4.3 that the minimal transversals of orientation-reversing circuits are cycles of a specific homology (mod 2) in the dual t -map. This theorem implies that the elimination of a reducer does not alter the cardinality of a transversal of orientation-reversing circuits. The reducers are defined for orientable surfaces as well. We show with an example (fig. 4.6) that t -maps free of reducers in surfaces of higher connectivity (even the torus) can be very complicated.

Chapter 1

Cubic Graphs, Three Colors, and Maps

1.A Maps and a -maps

The graphs which we use may contain loops and multiple edges. The set of edges of graph G is denoted by eG ; its set of vertices is denoted by VG . The two vertices which are incident to an edge are called the *ends* of the edge. If the two ends coincide the edge is called a *loop*.

The *valency* (or *degree*) of a vertex v of graph G , $val_G(v)$, is the number of edges incident to v where loops are counted twice. A subset of edges in eG is called a *perfect matching* if it consists of a set of non-incident edges which covers every vertex of G . A graph G is called *cubic*, as usual, if $val_G(v) = 3$ for every vertex v in VG . A graph G is a *polygon* if it has at least one edge, is connected, and $val_G(v) = 2$ for all v in VG .

A map M is an ordered triple (C_M, v_M, f_M) where:

- (i) C_M is a finite cubic graph;
- (ii) v_M and f_M are disjoint perfect matchings in C_M , such that the subgraph of C_M induced by $v_M \cup f_M$ is a collection of polygons with four edges or *squares*.

From the above definition, it follows that C_M may contain double edges but not loops.

A third perfect matching in C_M is $eC_M - (v_M \cup f_M)$ and is denoted by a_M . The set of diagonals of the squares, denoted by z_M is a perfect matching in the complement of C_M .

The edges in v_M, f_M, a_M are called respectively v_M -edges, f_M -edges, a_M -edges.

In this section we show a natural 1-1 correspondence between the above objects and the algebraic maps introduced by Tutte in [25]. As we show, they are essentially the same; however, the definition of orientability in our model is simpler. In the next section we relate our maps with the usual topological model.

Tutte's paper is a permutation representation for graphs embedded in arbitrary surfaces. Here, we call it an algebraic map. In this section we do not give the topological interpretation of the objects.

An *algebraic map* A , or *a-map* A , is an ordered triple of permutations,

$$A = (R, \Theta, \Phi),$$

acting on a finite set B satisfying the following axioms:

$$(am1) \quad \Theta\Phi = \Phi\Theta, \quad \Theta^2 = \Phi^2 = \text{identity};$$

$$(am2) \quad x, \Theta x, \Phi x, \Theta\Phi x \text{ are different for all } x \in B;$$

$$(am3) \quad R\Theta = \Theta R^{-1};$$

$$(am4) \quad R^n(x) \neq \Theta x, \text{ for all integer } n \text{ and all } x \text{ in } B.$$

If, in addition, the following property is satisfied, A is a *connected a-map*:

$$(amS) \quad \langle R, \Theta, \Phi \rangle \text{ is transitive in } B,$$

where $\langle R, \Theta, \Phi \rangle$ represents the group of permutations generated by R, Θ , and Φ .

In [25] axiom (am5) is imposed. If it does not hold, the object there is called a pre-map. Paper [25] and an observation due to V.A. Liskovets [30, pg.14], which consists essentially in the replacement of R by $R\Theta$ in the above axioms, were the basis for the definition of map that we give.

We define a function ψ^a , from the set of maps into the set of a -maps as follows: given map $M = (C_M, V_M, f_M)$, the a -map $\psi^a(M)$, which we simplify to M^a , is $M^a = (a_M v_M, v_M, f_M)$ acting on the vertices of C_M . For this purpose, we identify the perfect matchings in C_M, V_M, f_M , and a_M as fixed-point-free involutions on VC_M in the natural way: if b is a perfect matching in (a_M, v_M, f_M) and $x \in VC_M$, then $b(x)$ is the other vertex in VC_M such that x and $b(x)$ are the ends of an edge in b .

(1.1) Proposition: ψ^a is a bijection from the set of maps onto the set of a -maps.

Proof: Initially we show that M^a is indeed an a -map, by verifying Tutte's axioms for it. The squares formed by the perfect matchings v_M and f_M , and the interpretation of these as fixed-point-free involutions account for (am1) and (am2). Axiom (am3) also follows,

$$\begin{aligned} (a_M \circ v_M) \circ v_M &= a_M \\ &= v_M \circ (v_M \circ a_M) \\ &= v_M \circ (a_M \circ v_M)^{-1} \end{aligned}$$

We prove (am4) for M^a indirectly, assuming that it fails. Since VC_M is finite we can restrict ourselves to non-negative exponents in (am4). Choose the pair (n, x) such that n is the smallest possible non-negative integer with (am4) failing. First we show that $n > 1$. If $n = 0$, then $x = v_M(x)$, contradicting the fact that v_M is a matching in C_M . If $n = 1$, then $a_M v_M(x) = v_M(x)$, or $a_M(y) = y$, with $y = v_M(x)$; this contradicts the fact that a_M is a

matching in C_M . Thus $n > 1$. We then have

$$(a_M, v_M)^n(x) = v_M(x);$$

equivalently

$$(a_M v_M)^{n-1}(a_M v_M(x)) = v_M(x),$$

or

$$v_M a_M (a_M v_M)^{n-1}(a_M v_M(x)) = v_M a_M v_M(x).$$

Hence, $(a_M v_M)^{n-2}(y) = v_M(y)$ where $y = a_M v_M(x)$. This contradicts the choice of (n, x) ; thus, (am4) holds for M^a . The proof that M^a is an a -map is complete.

To prove that ψ^a is a bijection, we define a function $\bar{\psi}^a$ from the set of a -maps into the set of maps such that, functions $\bar{\psi}^a$ and ψ^a are inverses. Given an a -map $A = (R, \Theta, \Phi)$ acting on B we define $\bar{\psi}^a(A)$ as $M = (C_M, v_M, f_M)$, where $VC_M = B$, $v_M = \Theta$, $f_M = \Phi$, and $a_M = R\Theta$. The edges of C_M are defined by v_M, f_M, a_M . It follows that C_M is a finite cubic graph. Axioms (am1) and (am2) imply that the subgraph of C_M induced by $v_M \cup f_M$ is a collection of disjoint squares. Therefore, $M = \bar{\psi}^a(A)$ is a map. Observe that $\bar{\psi}^a$ can be seen as just the construction of a Cayley color graph for the action of $(R\Theta, \Theta, \Phi)$ over B , according to (9); we drop the directions on the edges, replacing a pair of opposite oriented parallel edges by a single unoriented edge. This is possible without losing information because $R\Theta, \Theta$, and ψ are involutions. It is evident from the definitions of ψ^a and $\bar{\psi}^a$ that they are inverses. This concludes the proof of the proposition. ■

Observe that (amS) holds for an a -map $A = (R, \Theta, \Phi)$ iff C_M is connected, where $M = \bar{\psi}^a(A)$. In general, the orbits of $\langle R, \Theta, \phi \rangle$ acting over VC_M are the components of C_M .

The *components* of a map M are the components of C_M . M is *connected* if C_M has one component.

The $v_M f_M$ -polygons in C_M are called the *squares* of M , or the M -

squares. The set of M -squares is denoted by $SQ(M)$. The $v_M a_M$ -polygons in C_M are called the v -gons of M . The set of v -gons of M is denoted by $vG(M)$.

The $f_M a_M$ -polygons in C are called the f -gons of M .

The set of f -gons of M is denoted by $fG(M)$.

The $z_M a_M$ -polygons in $C_M \cup z_M$ are called the z -gons of M . The set of z -gons of M is denoted by $zG(M)$. The vertices of C_M are also called the *corners* of M .

The *Euler characteristic* of a map M is the integer

$$\chi(M) = |vG(M)| - |SQ(M)| + |fG(M)|.$$

The *connectivity* of a map M is the integer $\xi(M)$ defined as $2 - \chi(M)$.

A map M is called *orientable* if C_M is bipartite; otherwise it is called *non-orientable*. The *orientability* label of M , $OL(M)$, is defined as 1 if M is orientable and as -1 otherwise.

An a -map $A = (R, \Theta, \Phi)$ acting on B is *non-orientable* (see [25]) if $\langle R, \Theta\Phi \rangle$ has one orbit for some orbit of $\langle R, \Theta, \Phi \rangle$. The alternative (as we show in the proof of the next proposition) is for $\langle R, \Theta\Phi \rangle$ to have two orbits for every orbit of $\langle R, \Theta, \Phi \rangle$. In this case, A is called orientable.

The concepts of orientability for maps and a -maps (as well as for topological maps, as we show later) agree.

We have the following proposition:

(1.2) Proposition: Map $M = (C_M, v_M, f_M)$ is orientable iff a -map M^a is orientable.

Proof: We can, evidently, restrict the proof to one component of C_M , corresponding to one orbit of $\langle a_M v_M, v_M, f_M \rangle$, acting on VC_M . In other

words, we may suppose C_M connected.

Rephrasing the proposition according to the definitions, we want to prove that C_M is bipartite iff $\langle a_M v_M, v_M f_M \rangle$ has more than one orbit on VC_M . Observe that the actions of $a_M v_M$ and $v_M f_M$ correspond to paths of length two in C_M . It follows that if C_M is bipartite, then $\langle a_M v_M, v_M f_M \rangle$ is not transitive on VC_M . To prove the converse, we establish as a lemma that there are at most two orbits of $\langle a_M v_M, v_M f_M \rangle$ acting on VC_M , for arbitrary connected M . To this end, we first show that all the four (or less) squares adjacent to a square S by a_M -edges are reachable by a path which starts in a fixed corner x in S , defined by a member of $\langle a_M v_M, v_M f_M \rangle$. Take the following four paths defined as

$$(i) (a_M v_M)(v_M f_M)x;$$

$$(ii) (v_M a_M)(v_M f_M)x;$$

$$(iii) (v_M a_M)x;$$

$$(iv) (a_M v_M)x.$$

Note that $v_M a_M = (a_M v_M)^{-1} \in \langle a_M v_M \rangle$ since VC_M is finite.

Observe that all the squares adjacent to S by an a_M -edge contain the final vertex of at least one of the above paths. By the connectivity of C_M , it follows that every orbit of $\langle a_M v_M, v_M f_M \rangle$ contains vertices of all squares. Since opposite vertices of a square are in the same orbit, it follows that there are at most two orbits of $\langle a_M v_M, v_M f_M \rangle$ on VC_M . Moreover, if there are two orbits, each contains two opposite vertices of each square.

Now, we can easily conclude the proof. If C_M is not bipartite, consider an odd polygon in C_M . At least two adjacent vertices, x and y , of this polygon are in the same orbit, as we proved that there are at most two orbits. We may consider that x and y are in the same square. If not, then they are linked by an a_M -edge and the replacement of y by $v_M a_M(y)$ fixes the situation. Therefore, we have two adjacent vertices of a square in the

same orbit of $\langle a_M v_M, v_M f_M \rangle$. Hence, by the argument above, which proves that each orbit contains two opposite vertices of each square, there is just one orbit. This concludes the proof. ■

With the above proposition we conclude the analysis relating a -maps and maps.

1.B Maps and t-maps

In this section we establish a natural 1-1 correspondence between our maps and the usual maps of graphs embedded on surfaces. Some properties of the latter objects are directly established in our model.

A *surface (without boundary)* is a Hausdorff topological space in which every point has a neighborhood homeomorphic to an open 2-ball, or open *disc*.

It is a classical theorem in topology that the surfaces which are connected, compact and without boundary are determined (up to homeomorphism) by the Euler characteristic and orientability.

The surfaces which we use have no boundary, are connected and, with the exception of the plane, are compact. In view of this we no longer use these qualifications. The plane, in reality, is simply a convenient way to replace the 2-sphere. Thus, we talk about planar maps where, to be precise, we should talk about spherical maps. We transform the sphere into the plane using a stereographic projection.

We denote the Euler characteristic of a surface S by $\chi(S)$. Also, let $OL(S)$ be $+1$ or -1 according to whether or not S is orientable.

Given a connected map M , we associate with M the surface S_M , characterized by $\chi(S_M) = \chi(M)$ and $OL(S_M) = OL(M)$.

In view of this, *planar* maps are connected maps M which have $\chi(M) = 2$ and C_M bipartite. We show in Theorem (2.0.2) that the latter condition about the bipartiteness of C_M is redundant.

Projective maps are connected maps M for which S_M is the surface known as the real projective plane. That means $\chi(M) = 1$ and C_M not bipartite. The latter condition is shown to be redundant in Theorem (1.7).

Toroidal maps are connected maps M for which $\chi(M) = 0$ and C_M is bipartite. S_M in this case is the familiar torus.

Kleinian maps are connected maps M for which $\chi(M) = 0$ and C_M is not bipartite. S_M in this case is the Klein bottle.

By a *path* in a graph G we mean a sequence $\pi = (v_0, e_1, e_2, \dots, v_{n-1}, e_n, v_n)$ such that $v_i \in VG$, $e_i \in eG$ and v_{i-1} and v_i are the ends of edge-term e_i for each i between 1 and n . A path is *degenerate* if it consists of just one vertex-term. Vertex v_0 is called the *origin* of π and vertex v_n is its *terminus*. The integer n is called the length of the path. A path is *reentrant* if its origin is equal to its terminus. A *cyclic* path is a reentrant path in which the sequence is considered cyclic, i.e., the origin can be adjusted to any vertex-term. If π is a path, the set of edges which appear in the sequence π as edge-terms is denoted by $e\pi$; analogously, the corresponding set of vertices is denoted by $V\pi$. If the terminus of π_1 is the same as the origin of π_2 , then we denote by $\pi_1 \circ \pi_2$ the composition of the paths π_1 and π_2 . The inverse of path π is denoted π^{-1} .

We proceed to show that the correspondence $M \rightarrow S_M$ is a natural one.

Assume that $\pi = \{\pi_1, \dots, \pi_f\}$ is a collection of cyclic paths in G that uses each one of its edges twice. Suppose also that the identifications of the ends of the edges in G forced only by the transitions in paths of π reproduce the vertices of G . Then π is called a *combinatorial embedding* of G .

For the next definition we consider a graph as a topological object, in the usual sense.

A graph G is *well-embedded* in a surface S if G is embedded in S , in the topological sense, and if the difference $S \setminus G$ is topologically equivalent to a collection of disjoint open discs.

A t -map X is a pair (G, S) in which graph G without isolated vertices is well-embedded in surface S .

Associated with a combinatorial embedding of a graph G , there is a natural t -map: let each one of the cyclic paths which defines the embedding bound a disc; identify pairwise the boundaries of these (disjoint) discs by “gluing” the two occurrences of each edge so as to identify correctly its ends (a loop is dealt with by the introduction of an artificial bivalent vertex), and so as to preserve the interior of each disc as an open disc.

The final object can be seen to form a t -map (G, S) , where S is a surface which has Euler characteristic $\chi(S) = v - e + f$; v, e , and f are, respectively, the number of vertices and edges of G , and the number of cyclic paths forming the combinatorial embedding. S is *orientable* if it is possible to choose a direction of traversal of the cyclic paths such that for each edge e with ends v and u , the traversal is once from v to u and once from u to v (a loop is, again, dealt with by the introduction of an artificial bivalent vertex).

The values of $\chi(S)$ and $OL(S)$, as explained above, are the same if we take an arbitrary graph H well-embedded in S . Therefore, the determination of $\chi(S)$ and $OL(S)$ is effected via some t -map (H, S) . The proof of this invariance is obtained by transforming an arbitrary t -map (H, S) into a canonical one, (H_S, S) , by means of elementary operations such as never to change $\chi(S)$ or $OL(S)$. This process is usually effected in connection with the classification of surfaces. (See Ringel [18] or Giblin [7]).

A dart is an oriented edge. The following proposition is straightforward. It characterizes orientability in two slightly different ways which are used in the proof of Theorem 1.6.

(1.3) Proposition: Suppose $\chi = (G, S)$ is a t -map. The following two

conditions are equivalent for the orientability of S :

- (i) Represent each facial path of χ by a polygon; let these polygons bound disjoint discs; S is orientable iff it is possible to simultaneously embed all these discs in the plane such that for each edge of G it is possible to “glue” along the two occurrences (in the boundary of the discs) of the edge “sliding” or “stretching” the discs, but not leaving the plane.
- (ii) Orient arbitrarily the edges of G ; as in (i), represent the facial paths of X by polygons and let these polygons bound disjoint discs; S is orientable iff it is possible to embed all the discs in the plane such that for each dart of G one occurrence in the boundary of the disc is clockwise and the other is counterclockwise.

For a t -map $X = (G, S)$ the disjoint open discs in $S \setminus G$ are called the *faces* of X . Observe that the boundary of each such disc corresponds in G to a cyclic path which we call a *facial path*. Note that an edge may occur twice in a facial path.

The (*combinatorial*) *boundary* of a face of X is the set of edges occurring once in the corresponding facial path. Observe that this notion agrees with the topological counterpart, and so, we drop the adjectives. A boundary in X is the mod 2 sum of the boundaries of faces. If F is a set of faces of X , the mod 2 sum of the boundaries of the faces in F is denoted by $\partial_x(F)$, or by $\partial(F)$ if X is understood.

Consider a cubic graph G with a proper 3-coloring of its edges in colors a, b, c . (This has the usual meaning of a partition of eG into three classes such that for each vertex the three edges incident to it are in different classes.)

The subgraph of G formed by edges of any two of the three colors consists of a number of disjoint polygons of even length, called ab -, bc -, ca -polygons.

The *faithful* embedding of G relative to a 3-coloring in colors a, b, c is given by the cyclic paths induced by the ab -, the bc -, and the ca -polygons. (It is evident that they form a combinatorial embedding.)

Given a map $M = (C_M, V_M, f_M)$ there is a natural 3-coloring of the edges of $C_M : \{v_M, f_M, a_M\}$. (Recall that a_M is the third perfect matching in C_M .)

The *embedding of a map* M is the faithful embedding of C_M relative to $\{v_M, f_M, a_M\}$.

Now we define a function ψ which is a bijection from the set of maps onto the set of t -maps. We denote $\psi(M)$ by M^t . Given a map M , to obtain M^t we proceed as follows. Consider the t -map $(C_M, S(M))$ given by the faithful embedding of M . The v -gons, the f -gons and the M -squares are boundaries of (closed, in this case) discs embedded (and forming) the surface $S(M)$. Shrink to a point the disjoint closed discs bounded by v -gons. The M -squares, then, become bounding digons. Shrink each such bounding digon to a line, maintaining unaffected its vertices. With these contractions, effected in $S(M)$, t -map $(C_M, S(M))$ becomes, by definition, $M^t = (G_M, S(M))$. Graph G_M is called the *graph induced by* M . A combinatorial description of G_M can be given as follows: the vertices of G_M are the v -gons of M ; its edges are the squares of M ; the two ends of an edge of G_M are the two v -gons (which may be the same) that contain the v_M -edges of the corresponding square.

It is evident that ψ is invertible: given a t -map we replace each edge by a bounding digon in its surface, and then expand each vertex to a disc in order to obtain the embedding of a map. Therefore, ψ^{-1} is well-defined; in fact, it is the dual of a useful construction in topology, namely, barycentric division. Thus, ψ is a bijection from the set of maps onto the set of t -maps.

It can be observed that ψ induces a bijection from the set of M -squares onto the set of edges of G_M . We frequently use this bijection to identify the sets $SQ(M)$ and eG_M .

A subset W of edges in a graph G is said to be a *cycle* in G if the subgraph induced by W has even valency at every vertex. If $W = \emptyset$, we have the null cycle. A circuit is a minimal non-null cycle. A circuit is, then, the set of edges of a polygon. The set of cycles of a graph G forms a vector space

under mod 2 sum. This space is denoted by $CS(G)$.

Bijection ψ induces for each map M a correspondence, denoted ψ_c^M , from the cycle space of $C_M, CS(C_M)$, onto the cycle space of $G_M, CS(G_M)$. The correspondence ψ_c^M is defined as follows: for $S \in CS(C_M)$, an edge $s \in eG_M$ is in $\psi_c^M(S)$ if square $s \in SQ(M)$ meets S in exactly one f_M -edge. With this definition, it is evident that $\psi_c^M(S)$ is a cycle in G_M and that ψ_c^M is surjective.

Every vector space that we use is over $GF(2)$. Therefore, we do not distinguish explicitly between a subset and its characteristic vector, nor between symmetric difference and mod 2 sum of characteristic vectors.

The following proposition permits the combinatorial definition of orientation-reversing circuits. An orientation-reversing circuit in a t -map M^t is a circuit in G_M which has in S_M a neighborhood homeomorphic to the Moebius strip.

(1.4) Proposition: ψ_c^M is a homomorphism, for every map M . Its kernel is the subspace of $CS(C_M)$ generated by the edges in the v -gons and in the squares of M .

Proof: Let S_1 and S_2 be cycles in C_M . We must show that $\psi_c^M(S_1 + S_2) = \psi_c^M(S_1) + \psi_c^M(S_2)$. An edge e of G_M is in $\psi_c^M(S_1 + S_2)$ iff exactly one f_M -edge of square $e \in SQ(M)$ is in $S_1 + S_2$. Therefore, $e \in \psi_c^M(S_1 + S_2)$ iff an even number of f_M -edges of square e belongs to one of S_1 and S_2 , and one f_M -edge of square e belongs to the other. The latter statement is equivalent to $e \in \psi_c^M(S_1) + \psi_c^M(S_2)$. This proves that ψ_c^M is homomorphism.

The image under ψ_c^M of the edges in an arbitrary square or v -gon of M is the null cycle in G_M . Thus, the space generated by the squares and v -gons is contained in $\text{Ker}(\psi_c^M)$. Conversely, suppose that $S \in \text{Ker}(\psi_c^M)$. The intersection of S with the edges of an arbitrary square has zero or two f_M -edges. Denote by T the cycle formed by the edges of the squares $Q \in SQ(M)$ such that $eQ \cap S$ contains the two f_M -edges of Q . It follows that $S + T$ has

no f_M -edges. Since $S + T$ is the edge set of a collection of polygons, (C_M is cubic), it follows that $S + T$ is formed by a collection of v -gons, whose edge-set is denoted by U . Hence, $S = T + U$, with T induced by squares, U induced by v -gons. Therefore, we conclude that $\text{Ker}(\psi_c^M)$ is contained in the space generated by the edge sets of squares and v -gons of M . The proof of the proposition is complete. $\blacksquare$

Since an element of the kernel of ψ_c^M has an even number of edges of C_M , it follows that if $\psi_c^M(S_1) = \psi_c^M(S_2)$, then $|S_1| \equiv |S_2| \pmod{2}$. This observation makes the following definition meaningful.

A cycle S in G_M is called an r -cycle in M^t if $\psi_c^M(S') = S$ and $|S'|$ is odd, for some cycle S' in C_M . If $|S'|$ is even and $\psi_c^M(S') = S$, then we say that S is an s -cycle in M^t .

We observe that the r -circuits in M^t are precisely the orientation-reversing circuits in M^t . This topological notion is not used; we work with our parity definition of r -cycle.

An r -loop in M^t is a loop in G_M which is the (only) edge of an r -circuit in M^t . The definition of s -loop is similar, replacing r -circuit by s -circuit.

Observe that a subset $T \subseteq eG_M$ is a boundary in M^t iff there exists a cycle T' of C_M , such that $\psi_c^M(T') = T$, and T' can be written as the mod 2 sum of some subsets of v -gons, f -gons, and squares.

For a map M , two cycles in G_M are *homologous* (mod 2) if their symmetric difference is a boundary in M^t . Homology (mod 2) is, thus, an equivalence relation.

The following proposition shows that S_M and $S(M)$ are the same surface.

(1.5) Proposition: For any map M we have

$$S_M = S(M).$$

Proof: First we prove that $\chi(S_M) = \chi(S(M))$. By definition $\chi(S_M) = \chi(M) = |vG(M)| - |SQ(M)| + |fG(M)|$. Since $(C_M, S(M))$ is a t -map, we use C_M to get $\chi(S(M))$. Graph C_M has $6|SQ(M)|$ edges and $4|SQ(M)|$ vertices. The number of cyclic paths that define the combinatorial embedding is $|SQ(M)| + |vG(M)| + |fG(M)|$. Using the definition of $\chi(S(M))$ via $(C_M, S(M))$ and simplifying we get

$$\chi(S(M)) - |VG(M)| - |SQ(M)| + |fG(M)| = \chi(S_M).$$

By Theorem 1.6, proved below, the faithful embedding of C_M relative to $\{v_M, f_M, a_M\}$ is orientable iff C_M is bipartite. Thus, Proposition (1.5) is complete, provided Theorem 1.6 is proved. ■

(1.6) Theorem: Assume we are given a cubic graph G and a proper 3-coloring of its edges. The faithful embedding of G relative to the 3-coloring defines an orientable surface if and only if G is bipartite.

Proof: Call the three colors a, b, c . Assume first that the faithful embedding is orientable. Consider disjoint discs which are bounded by the ab -, the bc -, and the ca -polygons. By Proposition 1.3 (i) it is possible to embed the discs in the plane, such that for an arbitrary edge we can “slide” one of the discs whose boundary contains one Occurrence of the edge, so as to “glue” it with its second occurrence. Note that they have correct orientation, without the need of “turning over” one of the discs. Therefore, after identification along the edge, it has an end in which the cyclic sequence of edges around it is of type $a - b - c$ clockwise, and an end where it is $a - b - c$ counterclockwise. Also, if two edges are incident to the same vertex, then this vertex is of type clockwise or counterclockwise for both edges. Thus, we can partition VG into clockwise and counterclockwise vertices. Moreover, each edge links a clockwise vertex to a counterclockwise one. It follows that G is bipartite.

Conversely, assume that G is bipartite. Orient the edges consistently from one class to the other, such that every vertex of one class is a source and every vertex in the other class is a sink. Embed the discs bounded by ab -polygons such that the a -darts are clockwise and consequently the b -darts are counterclockwise. Embed the discs bounded by bc -polygons such that the b -darts are clockwise and the c -darts are counterclockwise. Finally, embed the discs bounded by ca -polygons such that the c -darts are clockwise and the a -darts are counterclockwise. Since every dart appears once clockwise and once counterclockwise, it follows from Proposition 1.3 (ii) that the surface of the faithful embedding is orientable. ■

The next theorem establishes, using our definitions, the known fact that $\chi(M)$ is even for orientable maps. We need this result in Proposition 3.2.2.

(1.7) Theorem: Suppose that G is a cubic 3-edge-colored bipartite graph with $2k$ vertices. Then the parity of the number of faces of the faithful embedding of G relative to the 3-coloration is the same as the parity of k . Consequently, if M is an orientable map, then $\chi(M)$ is even.

Proof: Denote the colors by a, b , and c . Let (X, Y) be a bipartition of G . Since G is regular, $|X| = |Y| = k$. Consider the functions π_{ab}, π_{bc} , and π_{ca} acting on X , where $\pi_{ab}(u)$, for instance, is the terminus of the path of length 2 which starts at u , uses an a -edge and then a b -edge. It is clear that the image of u lies in X , by the bipartiteness of G . We prove that $\pi_{ab}, \pi_{bc}, \pi_{ca}$ are permutations acting on X . By the symmetry in the three colors, and since $|X|$ is finite, it is enough to show that π_{ab} is injective. Assume that $\pi_{ab}(p) = \pi_{ab}(q) = r$ for $p, q, r \in X$. Vertex r is the final vertex of the ab -paths which start at p and at q . But there is only one b -edge incident to r . Thus, this b -edge is used by both paths; we conclude that the other end of the a -edge incident to p is the same as the other end of the a -edge incident to q . Denote this common end by s (which is a vertex in Y). The fact that there is only one a -edge incident to s implies that $p = q$. We conclude that π_{ab} is injective, and thus a permutation acting on X .

Observe that the number of faces of the faithful embedding of G is the sum of the numbers of cycles of the permutations $\pi_{ab}, \pi_{bc}, \pi_{ca}$. Note also that the composition $\pi_{ab} \circ \pi_{bc} \circ \pi_{ca}$ is the identical permutation, hence an even permutation. It follows that the total number of even cycles of the three permutations is even. (Recall that an even cycle has odd parity, and vice-versa.) Thus, the parity of the number of cycles of π_{ab}, π_{bc} and π_{ca} is the same as the parity of the total number of their odd cycles. Make three copies of the vertices in X and use each copy to represent the cycles of one of the permutations, as polygons. The even polygons use an even number of the $3k$ vertices. Therefore, the parity of the total number of odd polygons is the same as the parity of $3k$, that is, the same parity as k . Since we proved that the parity of the number of faces in the faithful embedding is the same as the parity of the total number of odd cycles of the permutations π_{ab}, π_{bc} and π_{ca} , the first part of the theorem is established.

Suppose M is orientable. By definition, it follows that C_M is bipartite. Apply the first part of the theorem to the embedding of M . Since the number of vertices of C_M is a multiple of 4, it follows that the number of faces of the faithful embedding of C_M is even. To compute $\chi(M)$ we have to add

$4|SQ(M)| - 6|SQ(M)|$ to the number of faces, which we proved to be even. The proof is complete. $\blacksquare$

With this theorem we conclude the relations among Tutte's axioms [25], the classical topological notion of a graph well-embedded in a surface, and the definition that we use here. We showed that, under the bijections ψ^a and ψ , they are in fact equivalent.

The next theorem can be used to give an upper bound on the number of disjoint odd polygons which are subgraphs of C_M . We derive this consequence in Lemma 4.8. It is also used in Theorem 2.14 to give an upper bound on the maximum number of independent vertices in special graphs.

(1.8) Theorem: Assume that G is a 3-edge colored cubic graph. The mod 2 sum of the sets of edges in an arbitrary subset S of ab -, bc -, and ca -polygons, where a, b, c are the colors used to paint the edges of G , induces a disjoint set of polygons of even length.

Proof: Since G is cubic and the mod 2 sum of the edge-sets of polygons induces a cycle, it follows indeed that it induces a disjoint set of polygons. We just have to prove that each component induced by the mod 2 sum of bicolored polygons is a polygon with an even number of edges.

Consider an arbitrary such component, denoted K . Each edge in the polygon K is in an odd number of members of S , by definition of mod 2 sum. Observe that each edge of G appears in precisely two bicolored polygons. It follows that every edge of K appears in exactly one bicolored polygon in S . If an edge of K appears in an ab -polygon in S , let us mark it with $\bar{c}$. Analogously, mark an edge of K with $\bar{a}(\bar{b})$ if it appears in a bc -polygon (ca -polygon) in S . This partitions K into a number of maximal paths whose edges have the same mark, which we say is the label of the path. We observe that for x, y different symbols in $\{a, b, c\}$, if an edge of K marked $\bar{x}$ is incident to another marked $\bar{y}$, then the edge marked $\bar{x}$ is an y -edge and the edge marked $\bar{y}$ is an x -edge. From this observation we derive the following fact:

the number of edges in a maximal path with the same label is even if and only if the path that precedes it in K and the path that follows it have distinct labels.

The problem, then, is reduced to the following: if the vertices of a polygon (with at least three edges) are properly colored in at most three colors, then the number of vertices which have neighbors of the same color is even.

The vertices correspond to the maximal paths. Therefore, the vertices which have neighbors of the same color correspond to the maximal paths with an odd number of edges. If there are just one or two maximal paths, the theorem is trivial. We deal with the new problem in the next lemma.

(1.8.1) Lemma: Suppose the vertices of a polygon, with at least three vertices, are colored with at most three colors such that adjacent vertices have different colors. Then the number of vertices which have neighbors of the same color is even.

Proof: Let us denote the three colors by 0, 1, and 2. Orient the edges linking a vertex of color i to a vertex of color $i + 1 \pmod{3}$ from the former to the latter. It follows that the vertices which become a sink or a source with this orientation are precisely the vertices which have neighbors of the same color. The lemma follows if we observe that the number of sources is the same as the number of sinks. ■

The idea to orient the edges, which makes very simple the above proof, is due to H. Shank in a personal communication.

1.C The Phial Map and the Antimap

In this section we use our definition of map to derive a richer structure, which is the basis of the next chapter.

Given a map $M = (C_M, V_M, f_M)$, let us denote by Q_M the 4-regular graph obtained from C_M by adding to its edges the 1-factor z_M in the complement of C_M , defined as the diagonals of the squares.

By leaving out one of the three disjoint perfect matchings v_m, f_M, z_M of Q_M and permuting the order of the other two, we get a set of six maps, denoted by $\Gamma(M)$, and named as follows:

SYMBOL	DEFINITION	NAME
M	$(Q_M \setminus z_M, v_M, f_M)$	map M
$D_M = D$	$(Q_M \setminus z_M, f_M, v_M)$	dual of M
$D_M^\sim = D^\sim$	$(Q_M \setminus v_M, f_M, z_M)$	antidual of M
$P_M = P$	$(Q_M \setminus v_M, z_M, f_M)$	phial of M
$P_M^\sim = p^\sim$	$(Q_M \setminus f_M, z_M, v_M)$	antiphial of M
M	$(Q_M \setminus f_M, V_M, z_M)$	antimap of M

Observe that if $X \in \Gamma(M)$, then $\Gamma(X) = \Gamma(M)$. Observe also that D^t (the t -map associated with D) corresponds to the usual dual t -map, which interchanges faces and vertices relative to M^t .

Consider a t -map M^t . We now define some important cyclic paths which are analogous to the facial paths. Starting at an arbitrary point of the surface near an edge we proceed (in the surface) Parallel to the edge, crossing it in its middle, and continuing to go parallel to the edge and in the same direction as before, but now in the other “side” of the edge. Near a vertex we turn (without crossing points of the graph) and repeat the process with the new edge. Iteration of the above rules ends when we get back to the initial edge in the initial “side”. Clearly, we reach this stage and we reach it before we reach a “side” of an edge already traversed. The edges which indicates the way and the vertices which make the transitions between two edges must be recorded such as to form a cyclic path in G_M . Such cyclic path is called a *zigzag* in M^t .

Observe that the zigzag paths form another combinatorial embedding of G_M , since each edge is used twice in the collection of zigzag paths. This embedding corresponds to $M^\sim$. We claim that M^t and $(M^\sim)^t$ share a duality

relation: their facial paths and zigzag paths are interchanged, while the (pairs of inverses) cyclic sequences of edges around the vertices of their graphs are the same. This claim can be easily verified in the maps M and $M^\sim$: their f -gons and z -gons are interchanged, while the v -gons are the same.

The z -gons of maps M and D are the same. This corresponds to the fact that the cyclic sequence of edges in the zigzag paths of M^t and D^t are the same.

M and P clarify a duality relation between M^t and P^t . The phial pair of maps M and P have v -gons and z -gons interchanged, while the same f -gons. This means that M^t and P^t have the cyclic sequence of edges around the vertices and the cyclic sequence of edges in the zigzag paths interchanged, while having the same cyclic sequence of edges in the facial paths.

The zigzag paths are studied by Shank [24], in the restricted case of planar t -maps. In this context he names these paths “left-right paths”. Paper [24] Influenced the definition of zigzag paths and the subsequent development of the theory presented in Chapters 2 and 3.

We show in Figures 1.1, 1.2, and 1.3 the elements of $\Gamma(M)$, where M is the t -map formed by a cube embedded in the plane. The dual t -map, D^t , is the familiar octahedron embedded in the plane. Observe that $S_{D^\sim} = S_P$, which is the non-orientable surface of connectivity 4, as shown by the non-orientable handles labelled A and B in Figure 1.2. Finally $S_{P^\sim} = S_{M^\sim}$, which is the familiar torus. The reader can observe the relations among the edges around the vertices, the edges in the facial paths, and the edges in the zigzag paths.




Figure 1.1




Figure 1.2




Figure 1.3

From the definitions of its members, there exists a natural bijective correspondence between the squares of the elements of $\Gamma(M)$.

Two maps $M = (C_M, v_M, f_M)$ and $N = (C_N, v_N, f_N)$ are *isomorphic* if there exists a bijection b from VC_N onto VC_M such that for every $x \in VC_M$,

$$bv_M(x) = v_Nb(x)$$

$$bf_M(x) = f_Nb(x)$$

$$ba_M(x) = a_Nb(x)$$

Two members of $\Gamma(M)$ are said to be *square-identically isomorphic* if they are isomorphic and the isomorphism preserves the vertices of each square.

We use the structure of $\Gamma(M)$ to prove that the graphs which arise by doubling every edge of another graph can be embedded so as to form a self-dual t -map in a strong sense, which we now describe. Denote by G^d the graph obtained from a graph G by doubling every edge of G . Consider an arbitrary t -map E^t such that $G_E = G$. Denote by D^t the t -map obtained from E^t by replacing every edge of G by a bounding digon. Let M denote the dual of D . Consider the pair of dual maps formed by the phial P of M and the antidual

$D^\sim$ of M . Under these conditions, we show in the next theorem that P and $D^\sim$ are square-identically isomorphic. It is an easy property, which we prove in 2.1, that $G_D \cong G_{D^\sim}$ for every map D . Consequently, $G^d \cong G_D \cong G_{D^\sim}$ is embedded such as to be *edge-identically* isomorphic, i.e., the isomorphism between G_P and $G_{D^\sim}$ induces the same 1-1 correspondence between eG_P and $eG_{D^\sim}$ as the one given by the topological duality.

(1.9) Theorem: Let D^t be obtained from E^t by doubling each edge of G_E so as to form a bounding digon in S_E . Let M be the dual of D , P be the phial of M , and $D^\sim$ be the antimap of D . Then the pair of dual maps, P and $D^\sim$ are square-identically isomorphic.

Proof: It is sufficient to prove that M and $M^\sim$ are square-identically isomorphic. This is so because P is the phial of M and $D^\sim$ is the phial of $M^\sim$.

Denote by N the dual of E . Observe that G_M is equal to G_N with every edge subdivides by a bivalent vertex, called a *middle* vertex of G_M .

It is a straightforward property of a map M , whose graph G_M is obtainable by subdividing every edge of another graph by a middle vertex, that M and $M^\sim$ are square-identically isomorphic. To verify this, recall that the antimap of a map M replaces, in each square of M , the two f_M edges by its diagonals. (See Figure 1.4). To be convinced that the map and its antimap under the hypothesis are square-identically isomorphic, just twist the v -gons with four edges which correspond to the middle vertices. ■




Figure 1.4

Chapter 2

Rich Maps and Graphs

2.A Algebraic Preliminaries

Initially we state the facts from linear algebra that we use.

Consider a finite dimensional vector space W over a field K and a basis $\{a_i\}_{i=1,\dots,n}$ of W . Let the symmetric bilinear function

$$\langle \cdot, \cdot \rangle : W \times W \rightarrow K$$

be defined by $\langle u, v \rangle = \sum_{i=1}^n \lambda_i \gamma_i$, where $\lambda_i, \gamma_i \in K$, for $i \in \{1, \dots, n\}$, and $u = \sum_{i=1}^n \lambda_i a_i$, $v = \sum_{i=1}^n \gamma_i a_i$. Function $\langle \cdot, \cdot \rangle$ is the well-known scalar product induced by the basis $\{a_i\}_{i=1,\dots,n}$. Relative to the scalar product, there is the concept of orthogonality: two vectors $u, v \in W$ are *orthogonal* if $\langle u, v \rangle = 0$. If A is a subspace of W , then the set of vectors of W which are orthogonal to all vectors in A is also a subspace of W , denoted by $A^\perp$.

The following is a list of elementary algebraic properties, involving the above concepts. Proofs of these properties can be found in Godement's book [8, Sections 19, 36].

(2.0.1) For a finite dimensional vector space W , over an arbitrary field, let A and B denote arbitrary subspaces of W , and let “ $\perp$ ” be defined relative

to a scalar product induced by a fixed basis for W . The following properties hold:

- (a) $\dim(A + B) = \dim(A) + \dim(B) - \dim(A \cap B)$;
- (b) $(A^\perp)^\perp = A$;
- (c) $(A + B)^\perp = A^\perp \cap B^\perp$; $(A \cap B)^\perp = A^\perp + B^\perp$;
- (d) $\dim(A) + \dim(A^\perp) = \dim W$. ■

We also use the fundamental homomorphism theorem (for vector spaces): assume that W_1 and W_2 are vector spaces over the same field, and $h : W_1 \rightarrow W_2$ is a homomorphism. Then, the quotient space $W_1/\text{Ker}(h)$ is isomorphic to the subspace of W_2 defined by image of W_1 under h . In particular,

$$\dim[\text{Ker}(h)] + \dim[\text{Im}(h)] = \dim W_1.$$
■

If W is a subset of vertices of a graph G , the *coboundary* of W , denoted by $\delta_G(W)$ or $\delta(W)$, is the set of edges of G which have one end in W and the other in $VG - W$.

Let M be a given map. Consider the graphs G_M, G_D , and G_P . The sets of coboundaries of these graphs are denoted by V_M, F_M , and Z_M respectively. To simplify the notation, we use V, F , and Z , if M is understood.

We give, as usual, a vector structure to the power set $2^{SQ(M)}$ by taking the sum of subsets to mean their mod 2 sum.

Thus, the underlying field is $GF(2)$. Let the canonical basis for $2^{SQ(M)}$ be the set of singletons representing the individual squares of M .

With the choice of this basis we have a scalar product and the notion of perpendicular subsets of squares of M , (or subsets of edges of G_M, G_D or G_P) in this special case two subsets are *perpendicular* if their intersection has even cardinality.

Since the symmetric difference of two coboundaries in a graph G is also a coboundary in G , the set of coboundaries of G , denoted $BS(G)$, is a subspace of $2^{SQ(M)}$. Analogously, since the symmetric difference of two cycles in G is also a cycle in G , the set of cycles of G , denoted as said before by $CS(G)$, is a subspace of $2^{SQ(M)}$. For every graph G the intersection of any member of $BS(G)$ and any member of $CS(G)$ has even cardinality. Indeed, $BS(G)$ and $CS(G)$ are perpendicular subspaces, for every graph G . See Bondy and Murty [1]. The dimensions of these subspaces are

$$\begin{aligned}\dim[CS(G)] &= |eG| - |VG| + p \\ \dim[BS(G)] &= |VG| - p,\end{aligned}$$

where p is the number of components of G . See also [1].

In view of the above relation between $BS(G)$ and $CS(G)$, we denote by $V^\perp, F^\perp$ and $Z^\perp$, respectively, the cycles spaces of G_M, G_D , and G_P .

We use the dimensions of cycle and coboundary spaces of graphs and the theory of 3-colored cubic graphs to establish the familiar upper bound for $\chi(M)$, for connected maps M . We also show that if the upper bound is attained, then C_M is bipartite. This shows that in the definition of planar maps, the condition that C_M is bipartite is redundant.

(2.0.2) Theorem: The maximum Euler Characteristic of a connected map is 2. If $\chi(M) = 2$, then M is orientable.

Proof: We prove a more general statement. Suppose G is a connected cubic graph and $\{a, b, c\}$ is the set of colors of a proper 3-coloration of the edges of G . Let $2k$ be the number of vertices of G ; consequently, $3k$ be the number of its edges. Denote by $p(x, y)$ the number of xy -polygons, $x, y \in \{a, b, c\}$. We prove that $2k + p(a, b) + p(b, c) + p(c, a) - 3k$ is at most two, and that if this bound is attained, G is bipartite. Specializing for the faithful embedding of C_M , we have the theorem.

Observe that the dimension of the cycle space of G is $3k - 2k + 1 = k + 1$.

The dimension of the subspace of $CS(G)$ generated by the $ab-$, $bc-$, and $ca-$ polygons is $p(a, b) + p(b, c) + p(c, a) - 1$. To see this, observe that this subspace is the space of the coboundaries of the topological dual of the t -map formed by the faithful embedding of G , and that this dual has its vertices in bijective correspondence with the $ab-$, the $bc-$, and the $ca-$ polygons. Therefore, we have

$$p(a, b) + p(b, c) + p(c, a) - 1 \leq k + 1,$$

that is,

$$p(a, b) + p(b, c) + p(c, a) - k \leq 2,$$

as we stated. If we have equality, then the three types of polygons generate $CS(G)$. As every such generator has an even number of edges, it follows that G is bipartite. This concludes the proof. ■

The intersection of $BS(G)$ and $CS(G)$, subspace of 2^{eG} over $GF(2)$, has been named the of G (see [21]).

The dimension of the bicycle space of a graph is an important parameter in the theory that follows.

2.B A Ternary Relation Involving Cycles and Coboundaries

In this section we introduce the concept of rich map. A rich map induces three graphs on the same set of edges having the property that an arbitrary cycle in one of them is expressible as the symmetric difference of two coboundaries, one in each of the two other graphs.

(2.1) Proposition: For any map M we have $G \cong G_{M^\sim}$.

Proof: The M -squares, which are the edges of G_M are in natural 1-1 correspondence with the $M^\sim$ -squares which are the edges of $G_{M^\sim}$; namely,

to M -square S corresponds the $M^\sim$ -square $S^\sim$ such that $eS \cup eS^\sim$ induces in Q_M a subgraph isomorphic to K_4 . The vertices of G_M are the v -gons of M , which are the v -gons of $G_{M^\sim}$. Therefore, the sets of vertices of G_M and of $G_{M^\sim}$ are equal. Also, the ends of an M -square (i.e., an edge of G_M) are the same v -gons which are the ends of the corresponding $M^\sim$ -square (corresponding edge of $G_{M^\sim}$). This concludes the proof of the proposition.

■

(2.2) Proposition: For every map M , the cycle space of one of G_M, G_D, G_P contains the sum of the coboundary spaces (over $GF(2)$) of the other two.

Proof: By the symmetry in the three graphs given by the implication $X \in \Gamma(M) \Rightarrow \Gamma(X) = \Gamma(M)$ and by the last proposition we simply have to prove that $V^\perp \supseteq F + Z$. Indeed, the proof for arbitrary map M that $V^\perp \supseteq F$ suffices, because for $M^\sim$, $V_{M^\sim}^\perp = V^\perp$ and $F_{M^\sim} = Z$.

To prove that $F \subseteq V^\perp$, for arbitrary map M , we show that the elements of a set generating F are elements of $V^\perp$. To this aim, it is sufficient to prove that the set T of squares of M which have one f_M -edge in an arbitrary f -gon corresponds to a cycle in G_M . This is straightforward because T is the image of the f -gon under Ψ_c^M . The reason for the sufficiency is that T is a coboundary of the vertex in G_D which corresponds to (in fact, is) an f -gon. The proof of the proposition is complete. ■

The above proposition makes the following definition meaningful. The *(cycle) deficiency* of a map M is the dimension of the quotient space $V^\perp / (F + Z)$. A map is called *(cycle) rich* if

$$\text{def}(M) = \dim \left(\frac{V^\perp}{F + Z} \right) = 0.$$




Figure 2.1

The three graphs presents in Figure 2.1 are G_M , G_D , and G_P for a rich map M . A cycle in one of the graphs can be expressed as the mod 2 sum of coboundaries in the other two. We exemplify this relation as follows:

- (1) the circuit $\{5, 6, 8\}$ in the first graph can be written as

$$\delta(\{b, d\}) + \delta(\{\ell\}) = \{1, 2, 3, 4, 6, 8\} + \{1, 2, 3, 4, 5\} = \{5, 6, 8\};$$

- (ii) the circuit $\{2, 6, 7\}$ in the second graph can be written as

$$\delta(\{f, i\}) + \delta(\ell) = \{1, 3, 4, 5, 6, 7\} + \{1, 2, 3, 4, 5\} = \{2, 6, 7\};$$

- (iii) the circuit $\{6\}$ in the third graph can be written as

$$\delta(\{f, h, i\}) + \delta(\{a, c, d\}) = \{1, 2, 4, 5, 7, 10\} + \{1, 2, 4, 5, 6, 7, 10\} = \{6\}.$$

The explanation for this ternary symmetric relation is given by the next

theorem.

(2.3) Theorem. (Cycle-coboundary symmetric ternary relation): If M is a rich map, then the three graphs G_M, G_D , and G_P satisfy the following property: the cycle space (over $GF(2)$) of one of them is equal to the sum of the coboundary spaces of the other two.

Proof: Assume the Lemma 2.4 on equality of the deficiencies. The above theorem is a direct corollary of this lemma with zero as the common deficiencies of M, D , and P . ■

(2.4) Lemma. (Equality of deficiencies): For a map M , every two members of $\Gamma(M)$ have the same deficiency.

Proof: By the Absorption Property, proved as Theorem 2.5 below, applied to P in place of M , it follows that $V_P \cap F_P \subseteq Z_P$. But $V_P = Z_M = Z$, $F_P = F_M = F$, $Z_P = V_M = V$. Thus, $Z \cap F \subseteq V$ or $F \cap Z = V \cap F \cap Z$.


$$\begin{aligned} \text{def } M &= \dim \left(\frac{V^\perp}{F + Z} \right) = \dim V^\perp - \dim(F + Z) \\ &= (e - v + p) - (f - p) - (z - p) + \gamma, \end{aligned}$$

where e is the number of squares of M ; v, f, z are, respectively, the numbers of vertices of G_M, G_D, G_P ; p is the number of components of M ; and γ is the dimension of $V \cap F \cap Z$.

The above expression can be rewritten as

$$\text{def}(M) - (e + 3p + \gamma) - (v + f + z),$$

which is evidently symmetric in v, f, z . Since every element of $\Gamma(M)$ only induces a permutation of v, f, z in the above expression, the lemma follows, provided the Absorption Property (Theorem 2.5) is proved. ■



Dependence Structure for Theorem 2.3

The next theorem is the key for the symmetry in this theory.

(2.5) Theorem. (Absorption Property): For arbitrary map M we have $V \cap F \subseteq Z$.

Proof: Take an element $b \in V \cap F$. It follows that there exist $S \subseteq vG(M)$ and $T \subseteq fG(M)$ such that b is the coboundary of S in graph G_M and the coboundary of T in graph G_D . Take the subgraph of C_M which consists of the v -gons in S , and denote it by C_S . Consider also the subgraph C_T , defined analogously by the f -gons in T . Denote by R the subgraph of C_M induced by the symmetric difference of eC_S and eC_T . We claim the following two facts about R :

(2.5.1) For every corner x of M , the a_M -edge incident to x is in eR if and only if the a_M -edge incident to $z_M(x)$ (the corner opposite to x in its square) is also in eR .

(2.5.2) A square of M has exactly one pair of opposite a_M -edges which are

incident to it in eR iff the square corresponds to an edge of G_M (and of G_D) which is in b .

Let us assume the two claims and conclude the proof of the Absorption Property. Following that we establish the claims.

By claim (2.5.1), graph R defines naturally a set U of v -gons in map $P = P_M$. Specifically, the subgraph C_U of C_P is formed from R by deleting every v_{M-} and f_{M-} edge of R and including a z_{M-} edge, whenever it links a pair of corners such that the a_M -edges which are incident to them are in eR . Thus, C_U is formed by a subset U of $VG(P)$. By claim 2.5.2 the coboundary of U in G_p is b . Hence $b \in Z$, concluding the proof of the theorem, subject to the proofs of the claims (2.5.1) and (2.5.2). ■

Proof of the claims (2.5.1) and (2.5.2) Take a square X of M . Consider that X has i v_M -edges in eC_S and j f_M -edges in eC_T . Both i and j are in $\{0, 1, 2\}$. Since $b \in V \cap F$, it follows that $|eX \cap (eC_T + eC_S)|$ is even for every square X of M . Hence, we have

$$(i, j) \in \{(0, 0), (1, 1), (2, 0), (0, 2)\}.$$

Claim 2.5.1 is directly checked for the four cases above. Thus, (2.5.1) follows, since X is arbitrary.

Square X corresponds to an edge in b iff $(i, j) = (1, 1)$. Direct observation shows that of the four cases the only one which implies that exactly one pair of opposite a_M -edges is included in eR is also $(i, j) = (1, 1)$. Hence, the proof of (2.5.2) is complete. ■

Observe that with the proof of these claims every statement used in the proof of Theorem 2.3 is established.

For a map M we define the *facial deficiency* of M as the dimension of $V^\perp/F$.

(2.6) Proposition: The facial deficiency of a connected map M is $\xi(M)$.

Proof: The proof follows from the definition:

$$\begin{aligned} \dim \left(\frac{V^\perp}{F} \right) &= \dim(V^\perp) - \dim(F) \\ &= (e - v + 1) - (f - 1) \\ &= (2 + e) - (v + f) \\ &= 2 - \chi(M) = \xi(M), \end{aligned}$$

where $e = |SQ(M)|$, $v = |vG(M)|$, $f = |fG(M)|$. ■

Part (b) of the following corollary appears in Shank [24], proved by more specific techniques.

Corollary of (2.6) (a) Planar maps are rich; (b) the bicycle space of a planar graph G is generated by the cycles induced by the zigzag paths of M^t , where $G_M = G$.

Proof: That a planar map M is rich follows from the fact that $\xi(M) = 0$, which implies, by the previous proposition that $V^\perp = F$, or $V^\perp = F + Z$.

To establish (b) it is sufficient to show that $V \cap V^\perp = Z$. By part (a) and Lemma 2.4, it follows that $V + F = Z^\perp$. Replacing F by $V^\perp$ and taking the perpendiculars on both sides of the equality yields $V \cap V^\perp = Z$, by 2.0.1(c). This concludes the proof of the corollary. ■

Planar maps are simple examples of rich maps, simple in the sense that we do not need Z to generate $V^\perp$. What we propose to do now is to characterize, in a topological way, rich maps. With the characterization it is easy to get examples of rich maps in arbitrary surfaces.

Motivated by Theorem 2.3, concerning the symmetric ternary relation in cycles and coboundaries, we introduce the following definition.

A graph is called (*cycle*) *rich* if there are two other graphs with the same

set of edges as the first graph such that a cycle in one of the three graphs corresponds to the symmetric difference of two coboundaries, one in each of the other two graphs.

We define also the *map-deficiency* of a graph G to be $\min_M \{\text{def}(M) : G = G_M\}$. A graph G is called *map-rich* if its map-deficiency is 0, that is, if there exists a rich map M with $G_M = G$.

Theorem 2.3 implies that all map-rich graphs are rich graphs. Having no counter-examples so far, we conjecture the converse:

Conjecture: All rich graphs are map-rich graphs.

This conjecture if true, would complete the analogy between rich graph and planar graph.

Since we have not made any progress to settle or disprove this conjecture, we consider the easier problem for the characterization of map-rich graphs.

2.C Surface Characterizations of Map-Rich Graphs

This section establishes two characterizations (Theorem 2.9) of map-rich graphs. One of these is used to give a topological “recipe” for all map-rich graphs. The other is used to prove that there are graphs which are not map-rich.

A *medial map* is a t -map $X = (H, S)$ such that H is 4-regular and the faces of X are partitioned into two ordered classes called “black” and “white” such that every edge of H is incident to faces of different colors.

We define a function acting on the set of maps with image in the set of t -maps: $\psi'(M)$ is the t -map obtained from the embedding of M by contracting to a point in S_M each disc bounded by a square of M . We use the notation $\psi'(M) = M' = (H_M, S_M)$. The next lemma shows that this notation is

convenient.

(2.7) Lemma: The function ψ' is a bijection from the set of maps onto the set of medial maps.

Proof: First we prove that $(H_M, S_M) = M'$ is a medial map for every map M . It is evident that H_M is 4-regular. To prove that M' is properly 2-face-colorable, it is sufficient to show that (C_M, S_M) is 3-face-colorable; this is straightforward: color with black the discs bounded by v -gons, with white the discs bounded by f -gons, and with red the discs bounded by the squares of M . When we contract the squares to get M' , the red faces disappear and we have the ordered 2-coloring, black and white, of the faces of M' . This shows that M' is a medial map.

Conversely, we show that $(\psi')^{-1}$ is defined for all medial maps. Given a medial map $X = (H, S)$, expand in S the 4-valent vertices of H to discs bounded by squares. Of course, this expansion gives the embedding of a map, denoted M : the original edges of H are the a_M -edges; the created edges which bound a black face are the v_M -edges; the created edges which bound a white face are the f_M -edges. It is evident that the v -gons, the f -gons, and the squares of M are boundaries of closed discs in S . The lemma is proved. ■

Observe that if $X, Y \in \Gamma(M)$, then $H_X = H_Y$. In the proof of the above lemma, the black faces of M' correspond to the v -gons of M , the white faces to the f -gons. Henceforth, we fix this convention.

Note that the only difference between M' and D' is that the black and white faces are interchanged.

The set of black faces of M' is denoted by $bF(M')$; its set of white faces is denoted by $wF(M')$.

Given a map M and a vertex v of valency $2n$ in G_M , two edges incident to v are opposite at v if their indices differ by n in a successive numbering of

the edges incident to v that follows the cyclic sequence induced by M^t .

For an eulerian G_M , a *smooth path* in M^t is a path in G_M such that every pair of consecutive edges is opposite at the vertex between them.

Observe that for G_M eulerian, the smooth paths of M^t define a partition of eG_M . In particular the smooth paths of $M' = (H_M, S_M)$ define a partition of eH_M .

Given M' we define G'_M as follows: the vertices of G'_M are the smooth paths of M' . The edges of G'_M are the vertices of H_M . The two ends of an edge of G'_M are the two smooth paths (which may be the same) that use the corresponding vertex of H_M .

(2.8) Lemma: For every map M , $G_P \cong G'_M$, where P is the phial of M .

Proof: Consider $M = (H_M, S_M)$. There exists a natural bijection, denoted b_1 , from the squares of P onto the vertices of H_M : the expansion in S_M of the vertices of H_M gives the squares of M , which are identifiable with the squares of P . Also there is a natural bijection from the v -gons of P onto the smooth paths of M' ; we denote by b_2 this bijection.

Take an edge x of G_P and let u and w be its ends. It follows, from the definitions, that the ends of $b_1(x) \in eG'_M$ are $b_2(u)$ and $b_2(w)$, which are in VG'_M . Therefore, $G_P \cong G'_M$, proving the lemma. ■

Given a map M we say that P' , the medial of the phial of M , is a *smooth-path-embedding* of G_M , or an *sp-embedding* of G_M . The *surface of the sp-embedding* P' is S_p . A graph is *eulerian* if every vertex has even valency.

(2.9) Theorem. (Surface characterizations of map-rich graphs):

- (a) The following three conditions are equivalent for connected graph G with at least one edge:

- (i) G is a map-rich graph.
 - (ii) G has an sp -embedding in a surface S of Euler characteristic $\chi(S) = 3 - |VG| + \gamma$, where γ is the dimension of the bicycle space of G . (The value of $OL(S)$ is also determined; see part (b)).
 - (iii) There exists an sp -embedding $P' = (H_P, S_P)$ of G such that the cycle space of H_P (over $GF(2)$) is generated by the boundaries of the faces of P' and by the cycles induced by the smooth paths of P' .
- (b) For every pair of phial maps, M and P , if P is orientable, then G_M is eulerian. The converse also holds if the maps are rich.

Proof of part (a) Initially we prove that (i) implies (ii). Assume that G is a map-rich graph and that M is a rich map satisfying $G_M = G$. Denote by P , as usual, the phial of M . By the lemma on bicycle spaces, proved below in 2.9.2,

$$Z \cap F = V \cap F \cap Z = V \cap V^\perp$$

where V, F , and Z are relative to map M . We have also

$$V = F^\perp \cap Z^\perp,$$

since M is rich and thus $V^\perp = F + Z$. We use the two latter equalities in the expression for $\xi(P)$ which we obtain from Lemma 2.9.1, proved below, with P replacing M :

$$\begin{aligned} \xi(P) &= \dim(Z' \cap F^\perp) - \dim(Z \cap F) \\ &= \dim(V) - \dim(V \cap V^\perp). \end{aligned}$$

Since G is connected, $\xi(P) = 2 - \chi(S_P)$, and $\dim(V \cap V^\perp) = \gamma$ it follows that

$$\xi(S_P) = 3 - |VG| + \gamma.$$

Therefore, since S_P is the surface of the sp -embedding of G , the proof that (i) implies (ii) is complete provided Lemmas 2.9.1 and 2.9.2 are established.

Now we prove that (ii) implies (i). Assume that G has an sp -embedding P' such that $\xi(S_p) = 3 - |VG| + \gamma$. Since G is connected the latter equality is equivalent to

$$\xi(P) - \dim(V) - \dim(V \cap V^\perp),$$

where V is the coboundary space of G_M , and M the phial of P . By Lemma 2.9.1, with P in the role of M ,

$$Z \cap F \subseteq V \cap V^\perp \subseteq V \subseteq Z^\perp \cap F^\perp$$

$$\dim(Z^\perp \cap F^\perp) - \dim(V \cap F) = \xi(P).$$

The previous expression for $\xi(P)$ and the above two properties imply


$$Z \cap F = V \cap V^\perp$$

$$V = Z^\perp \cap F^\perp.$$

Taking perpendicular to both sides of the latter equality we get

$$V^\perp = F + Z,$$

whence M is rich. Since by Lemma 2.8, $G_M = G'_P$; and the latter is G , it follows that G is map-rich. Thus, the proof that (ii) Implies (i) is complete provided Lemma 2.9.1 is established.



Dependence Structure for Theorem 2.9

We show that (i) and (iii) are equivalent. To this end, we define the *sp-deficiency* of a connected map P , or $\text{def}'(P)$, as follows. Denote the cycle space of H_p by $CS(H_p)$. Let the subspaces of $CS(H_p)$ generated by the boundaries of the black and white faces of P' be denoted, respectively, by Z' and F' ; denote by V' the subspace of $CS(H_p)$ generated by the cycles induced by the smooth paths of P' . With this terminology we have, by definition,

$$\text{def}'(P) = \dim \frac{CS(H_p)}{V' + F' + Z'}.$$

By Lemma 2.9.3, proved below, it follows that $\dim[(F' + Z') \cap V'] = 1 + \dim(F \cap Z)$.

We also have $\dim[CS(H_p)] = 2e - e + l = e + 1$, and $\dim(F' + Z') = f + z - l$, where $f = |vG(D)|$, $z = |vG(P)|$, $e = |eG_p|$; the first of those equalities is implied by the fact that P is connected; the second follows from the facts that $\dim(F') = f$ and $\dim(Z') = z$, because the boundaries of the black faces and the boundaries of the white faces of P' partition eH_p , and from the fact that $\dim(F' \cap Z') = 1$, proved in Proposition 2.9.4.

Since the edges in the smooth paths of P' also partition eH_p , it follows that $\dim(V)' = v$.

The above equalities imply that $\text{def}(P) = \text{def}'(P)$, for every map P :

$$\begin{aligned}
 \text{def}'(P) &= \dim[CS(H_p)] - \dim(V' + F' + Z') \\
 &= (e + 1) - \dim(F' + Z') - \dim(V') + \dim[(F' + Z') \cap V'] \\
 &= (e + 1) - (f + z - 1) - v + 1 + \dim(F \cap Z) \\
 &= e + 3 + \dim(F \cap Z) - (v + f + z) \\
 &= \text{def}(P).
 \end{aligned}$$

Now we can easily conclude the proof that (i) and (iii) are equivalent. If G is map-rich, let M be a rich map with $G_M = G$. By Lemma 2.0, this implies that $G'_p = G$. Since $\text{def}'(P) = \text{def}(P)$, as shown above, and $\text{def}(P) = \text{def}(M)$, by Lemma 2.4, the equality $\text{def}(M) = \text{def}'(P)$ shows that (iii) follows from (1).

Conversely, suppose that (iii) holds for some P' . Let M be the phial of P . Since $0 = \text{def}'(P) = \text{def}(P) = \text{def}(M)$, it follows that M is rich. By Lemma 2.8 we have $G_M = G'_p$; since $G'_p = G$, it follows that $G_M = G$ and so G is map-rich.

Thus, the proof of the equivalence between (i) and (iii) is reduced to the proof of Lemma 2.9.3 and Proposition 2.9.4.

Proof of part (b) Suppose that P is orientable, whence C_p is bipartite. Let $\{A, B\}$ be a bipartition of C_p . It follows, from the definition of phial, that for every pair of corners of M, X and $a_M(v_M(X))$, one is in A and the other is in B . Thus, the number of vertices in every v -gon of M is a multiple of four. Since the valency of a vertex in G_M is half the number of vertices in the corresponding v -gon of M , it follows that G_M is eulerian.

To prove the converse assume that M is rich and C_M is eulerian. (If we drop the hypothesis that M is rich, then the converse does not follow in general.) We use the equivalence between (i) and (iii) established in part (a). Consider the graph G of part (a) to be G_M . Therefore, (iii) holds since M

is rich. Let P be the phial of M . By (iii), we have $CS(H_p) = Z' + F' + V'$, where to recall the terminology, Z' and F' are generated by the boundaries of, respectively, the black and the white faces of P' , and V' is generated by the cycles induced by the smooth paths of P' .

Since G_M is eulerian it follows that each smooth path in P' has an even number of edges. This claim is justified as follows. Assume that there are k_1 loops and k_2 non-loops incident to a vertex v of $G = G_M$. The subgraph of H_p induced by the corresponding smooth path in P' has k_1 4-valent vertices and k_2 bivalent vertices. The number of edges in this subgraph is, therefore, $k = \frac{1}{2}(4k_1 + 2k_2)$. Since G_M is eulerian k_2 is even and so is k .

To each smooth path in P' corresponds a pair of (v_p, f_p, a_p) -paths (a cyclic path that uses v_p, f_p , and a a_p -edges in this order) in C_p such that the symmetric difference of their edge-sets is the edges in the squares that they meet once. Observe that since each smooth path has an even number of edges, so does each (v_p, f_p, a_p) -path.

We claim that the cycle space of C_P is generated by the cycles induced by the squares, the v -gons, the f -gons, and the (v_p, f_p, a_p) -paths. Observe that if the claim is accepted the proof is complete: as every generator has an even number of elements it follows that every cycle in C_p has an even number of edges, that is, C_p is bipartite and S_p is orientable. The claim is proved by the inclusion of a bounding (in S_P) loop at each vertex of H_p , and by the successive expansion of edges to form C_P . The proof uses the following two easy observations:

- (I) The cycle space of a graph G is generated by a collection X of cycles iff the cycle space of the graph G^e obtained from G by adjoining a new edge e which forms a loop, is generated by the collection X^e of cycles, obtained from X by including $\{e\}$ as one of its elements.
- (II) The cycle space of a graph G is generated by a collection X of cycles iff the cycle space of the graph G_e , obtained from G by contracting non-loop e , is generated by the collection X_e of cycles, obtained from X by taking the difference of its elements with $\{e\}$.

It is evident that (I) and (II) hold. Iterated application of them transforms (H_p, X) into (C_p, Y) , where X is the set of cycles formed by the boundaries of the faces of P' and by the edges in the smooth paths of P' ; X generates $CS(H_p)$; Y is the set of cycles induced by the squares, the v -gons, the f -gons of P , and by one member of each pair of associated (v_p, f_p, a_p) -paths; Y generates $CS(C_p)$. The proof of Theorem 2.9 is complete provided statements 2.9.1 to 2.9.4 are established. ■

(2.9.1) Lemma. (Triple inclusion property): For every map M we have

$$V \cap F \subseteq Z \cap Z^\perp \subseteq Z \subseteq V^\perp \cap F^\perp$$

and

$$\dim(V^\perp \cap F^\perp) - \dim(V \cap F) = \xi(M).$$

Proof: We restrict the proof, without loss of generality, to connected maps. To prove the equality, consider the following equation, with $e = |SQ(M)|$, $v = |vG(M)|$, $f = |fG(M)|$:

$$\dim(V^\perp \cap F^\perp) = (e - v + 1) + (e - f + 1) - \dim(V^\perp + F^\perp).$$

Using items (d) and (e) of 2.0.1 we have

$$\dim(V^\perp + F^\perp) = e - \dim[(V^\perp + F^\perp)^\perp] = e - \dim(V \cap F).$$

Using the above value of $\dim(V^\perp + F^\perp)$ yields

$$\dim(V^\perp \cap F^\perp) = (e + 2) - (v + f) + \dim(V \cap F) = \xi(M) + \dim(V \cap F),$$

which is the equality that we wish to establish.

To verify the inclusions we use the Absorption Property, Lemma 2.5, which says that

$$V \cap F \subseteq Z.$$

Proposition 2.2 implies that $Z^\perp \supseteq V + F$. Taking perpendiculars we get, by 2.0.1(c)

$$Z \subseteq V^\perp \cap F^\perp.$$

Thus, we have $V \cap F \subseteq Z \subseteq V^\perp \cap F^\perp$. To conclude the proof it is enough to show that $V \cap F \subseteq Z^\perp$. This follows from $V \cap F \subseteq V + F \subseteq Z^\perp$. The latter inclusion is implied by Proposition 2.2. The proof of the lemma is complete. ■

(2.9.2) Lemma on bicycle spaces: If M is a rich map, then the bicycle spaces of G_M, G_D , and G_p are equal to $V \cap F \cap Z$.

Proof: Observe that this lemma is a generalization of part (b) of the corollary of Proposition 2.6. Therefore, here is a property of planar graphs which generalizes to rich graphs.

The fact that M is rich implies that $V = F^\perp \cap Z^\perp$. Hence, $V \cap V^\perp = F^\perp \cap Z^\perp \cap V^\perp$. The symmetry in V, F, Z , shows that $F \cap F^\perp$ and $Z \cap Z^\perp$ are also equal to $V^\perp \cap F^\perp \cap Z^\perp$. We have $V \cap F \cap Z = V \cap F$, which follows from the Absorption Property (Lemma 2.5). Also we have $V \cap F \subseteq V \cap V^\perp$ by property 2.2. Since $V \cap V^\perp = V^\perp \cap F^\perp \cap Z^\perp$, it follows that $V \cap F \cap Z \subseteq V^\perp \cap F^\perp \cap Z^\perp$. To prove the reverse inclusion let $x \in V^\perp \cap F^\perp \cap Z^\perp$. As shown above the latter space is equal to $V \cap V^\perp = F \cap F^\perp = Z \cap Z^\perp$. Therefore, $x \in V \cap F \cap Z$ and $V^\perp \cap F^\perp \cap Z^\perp \subseteq V \cap F \cap Z$, concluding the proof. ■

(2.9.3) Lemma: For every connected map $P = P_M$,

$$\dim[(F' + Z') \cap V'] = 1 + \dim(F \cap Z).$$

Proof: We intend to prove the above equality by presenting an epimorphism h from $(F' + Z') \cap V'$ onto $F \cap Z$ and by showing that the kernel of h has dimension 1.

By Proposition 2.9.4, proved below, $F' \cap Z'$ has dimension 1 and hence two elements. It is also shown that the two elements of $F' \cap Z'$ are the null set of edges and the whole set of edges of H_p . Let $x \in F' + Z'$. By the preceding argument, there are two ways to represent x as mod 2 sum of boundaries of white and black faces of P' . Let W_x and B_x be subsets of $wF(P')$ and $bF(P')$ such that

$$x = \partial(W_x) + \partial(B_x).$$

Since the non-null element of $F' \cap Z'$ is eH_p , and $eH_p = \partial(wF(P')) = \partial(bF(P'))$, we have

$$\begin{aligned} x &= [\partial(w_x) + \partial(wF(P'))] + [\partial(B_x) + \partial(bF(P'))] \\ &= \partial(wF(P') \setminus W_x) + \partial(bF(P') \setminus B_x), \end{aligned}$$

which is the other way to represent x as mod 2 sum of boundaries of white and black faces of P' .

Note that if $x \in (F' + Z') \cap V'$, then for an arbitrary vertex v of H_p , the intersection of the edges incident to v with x consists of either zero, one, or two pairs of opposite edges. Moreover, this intersection is one pair if and only if v (which is an edge of G_D and G_p) is in the coboundaries of $W_x \subseteq VG_D$, and $B_x \subseteq VG_p$, in the adequate graphs. Therefore, $\delta_{G_D}(W_x) = \delta_{G_p}(B_x)$.

The function h from $(F' + Z') \cap V'$ into $F \cap Z$ is defined as follows. Given $x \in (F' + Z') \cap V'$, represent x as $\partial(W_x) + \partial(B_x)$ for convenient subsets of white and black faces. Take $h(x)$ to be the coboundary of W_x in graph G_D , which is equal, as explained above, to the coboundary of B_x in graph G_p . Since the coboundaries of a subset and its complement are the same, it follows that $h(x)$ is independent of the representation of x . Thus, h is well defined.

Now we show that h is a homomorphism. Take x and y in $(F' + Z') \cap V'$. Find W_x, B_x, W_y, B_y such that

$$x = \partial(W_x) + \partial(B_x),$$

$$y = \partial(W_y) + \partial(B_y).$$

Adding mod 2 we get

$$x + y = \partial(W_x + W_y) + \partial(B_x + B_y).$$

Thus,

$$h(x + y) = \partial(W_x + W_y) = \delta(W_x) + \delta(W_y) = h(x) + h(y).$$

Hence, h is a homomorphism. To prove that h is surjective take an element $u \in F \cap Z$. By definition, there exist $W_u \subseteq VG_D$ and $B_u \subseteq VG_p$ such that

$$u = \delta(W_u) = \delta(B_u).$$

Consider W_u and B_u as the subsets of the corresponding white and black faces of P' . Define $x \in F' + Z'$ as $x = \partial(W_u) + \partial(B_u)$. It can be directly observed that the intersection of x with the four edges incident to an arbitrary vertex consists of zero, one, or two pairs of opposite edges. Therefore, $x \in (F' + Z') \cap V'$. Moreover, $h(x)$ is the coboundary of W_u in G_D that is, $h(x) = u$. Thus, h is an epimorphism.

Now we prove that the kernel of h has dimension 1. Suppose that $h(x) = \phi$ and x is not the null vector. Find W_x and B_x such that $x = \partial(W_x) + \partial(B_x)$. One of W_x and B_x is non-empty, for instance, $B_x \neq \phi$. If some black face b is in B_x , then all the black faces adjacent to b by a vertex of H_p are also in B_x : where this not true, then $h(x) = \partial(B_x) \neq \phi$, contradicting the hypothesis. By the connectivity of P , it follows that $B_x = bF(P')$. The same argument shows that if $W_x \neq \phi$, then $W_x = wF(P')$. But if $B_x = bF(P')$ and $W_x = wF(P')$, then x is empty, contradicting the hypothesis. Thus, if $x \neq \phi$ and $h(x) = \phi$, then $x = \partial(bF(P')) = \partial(wF(P')) = eH_p$. Hence, the kernel of h has dimension 1. By the fundamental homomorphism theorem, it follows that

$$\dim[(F' + Z') \cap V'] = \dim[\text{Ker}(h)] + \dim[\text{im}(h)]$$

or

$$\dim[(F' + Z') \cap V'] = 1 + \dim(F \cap Z),$$

proving the lemma, subject to the establishment of Proposition 2.9.4, which we do next. ■

(2.9.4) Proposition: For connected $P = P_M$ the dimension of $F' \cap Z'$ is 1.

Proof: Assume that $x \in F' \cap Z'$. It follows that there are $W_x \subseteq wF(P')$ and $B_x \subseteq bF(P')$ such that $\partial(W_x) = x = \partial(B_x)$. If a face $b \in B_x$, then the elements of $wF(P')$ which are adjacent to b by an edge of H_p are in W_x . Therefore, by the connectivity of P , it follows that if both of W_x and B_x are not empty, then $W_x = wF(P')$ and $B_x = bF(P')$. Hence, if $x \in F' \cap Z'$ then x is either the null set or the whole set of edges, eH_p . Thus, $\dim(F' \cap Z') = 1$ as stated. ■

Observe that at this point every lemma and proposition used to prove Theorem 2.9 is established. To discover the surface with maximum Euler characteristic for which a (usual) embedding of a given graph exists is, in general, a difficult problem. In attacking the famous Heawood Conjecture, Ringel, Youngs, and others have determined the surface of maximum Euler characteristic where a (usual) embedding of a complete graph exists. See Ringel [18], Youngs [29]. With our next theorem we derive a natural upper bound on the Euler characteristic of a surface where there exists an sp -embedding of a given graph. This upper bound holds as an equality if and only if the graph is map-rich. In the next section we show that all complete graphs are map-rich.

(2.10) Theorem: Given a connected graph G and an sp -embedding of G , $P' = (H_p, S_p)$, then we have

$$\chi(S_p) \leq 3 - v + \gamma,$$

where γ is the dimension of the bicycle space of G , and $v = |VG|$. Moreover, equality holds iff P is rich.

Proof: The proof of this theorem is implicit in Theorem 2.9. However, we give a direct proof. Let M be the phial of P . Hence, G is G_M , by Lemma 2.8, and so G inherits the usual terminology for M . Since G is connected, we have

$$\begin{aligned} \text{def}(P) &= \dim Z - \dim(V + F) \\ &= (e - z + l) - (v - l) - (f - l) + \dim(V \cap F) \\ &= (e + 3) - (v + f + z) + \dim(v \cap F). \end{aligned}$$

Observe that $V \cap F \subseteq V \cap V^\perp$, by Proposition 2.2. Since $V \cap V^\perp$ is the bicycle space of G we have

$$0 \leq \text{def}(P) \leq (e + 2) - (f + z) - (v - 1) + \gamma,$$

which implies $\chi(S_p) \leq 3 - v + \gamma$, proving the inequality. if P is rich, then by the lemma on bicycle spaces, 2.9.2, and by the Absorption Property, 2.5, we have

$$V \cap V^\perp = V \cap F \cap Z = V \cap F.$$

Moreover, the value of $\text{def}(P)$ is 0. Thus, equality holds.

Assume that the equality holds, $\chi(S_p) = 3 - v + \gamma$. Then we have

$$f + z - e = 3 - v + \gamma,$$

or

$$0 = (e + 3) - (v + f + z) + \gamma.$$

Since $\gamma \leq \dim(V \cap F)$, by Proposition 2.2, it follows that $0 \leq \text{def}(P)$. But the definition of a map is at least zero. Hence, $\text{def}(P) = 0$, i.e., P is rich. ■

2.D Map-Rich and Non-Map-Rich Graphs

We use Theorem 2.9 to show that all complete graphs are map-rich, and to show that there are graphs which are not map-rich.

For complete graphs with an even number of vertices, K_{2n} , it is an easy matter to construct sp -embeddings in the real projective plane. To see this choose $4n$ points in the boundary of a disc, and link by chords opposite points, such that any two chords meet exactly once; if more than two chords meet at the same point, effect a local deformation of some of them, to make every crossing point the intersection of two chords. Now, identify every point in the boundary of the disc with its opposite, or antipodal point. With this antipodal identification the system of chords is now a t -map, with the real projective plane as its surface. The crossing points are the vertices of the graph, and its edges are the segments of chords between two crossings. By construction, this graph is 4-regular. We show that the t -map is properly 2-face-colorable. This is evident if $n = 1$. Assume that it is true for n at most k , and let n be $k + 1$. Observe that every pair of chords is the boundary of a subset R of faces. Remove two chords, c_1, c_2 , and obtain, by the inductive hypothesis, a proper 2-face-coloring of the remaining system, which has $2k$ chords. Put back the two chords. Observe that all the edges in the boundary of R , the set of faces bounded by c_1 , and c_2 , have the same color on both sides. Edges not in the boundary of R have different colors in each side. Therefore, to get a proper 2-coloring just interchange the colors in the faces of R . This completes the induction. Therefore, the projective t -map becomes after ordering the bicoloration, a medial map. A medial map that arises by the above construction is called a *simple $2n$ -system of projective lines*. Observe that if P' is such an object, then $G_M = G'_p = K_{2n}$.

(2.11) Theorem: For every n , the complete graph K_{2n} is map-rich.

Proof: To prove that K_{2n} is map-rich it is sufficient to show that if $P' = (H_p, S_p)$ is a simple $2n$ -system of projective lines, then M is rich: since in

this case $G_M - K_{2n}$, it follows that K_{2n} is map-rich. By Theorem 2.9 (iii) it is enough to show that the cycle space of H_p is generated by the boundaries of the faces of P' and the cycles induced by the smooth paths. Since S_p is the projective plane, the facial deficiency of P is 1, by Proposition 2.6. Therefore, we just need one cycle induced by a smooth path not spanned by the boundaries of the faces of P' . Observe that each one of these cycles is non-bounding, and therefore, none of them is in the span of the boundaries of the faces. This concludes the proof that every K_{2n} is map-rich. ■

We now want to extend the previous theorem to complete graphs with an odd number of vertices.

(2.12) Theorem: Assume that M is rich and $G_M = K_{2n}$. Then deleting in M^t all the edges incident to a vertex of K_{2n} , we have a rich map N with $G_N = K_{2n-1}$.

Proof: We use the labels $\{1, 2, \dots, 2n\}$ for the vertices of $G_M = K_{2n}$. Let $[i, j]$ denote the edge with ends i and j . We delete $[2, 2n], [3, 2n], \dots, [2n-1, 2n]$, in this order, obtaining, respectively, $M_2 M_3, \dots, M_{2n-1}$. Let F_i and Z_i be, respectively, the coboundary spaces of G_{D_i} and G_{P_i} for $i = 2, \dots, 2n-1$. By Proposition 2.12.1, proved below, $[i, 2n]$ is an edge of an element of $F_{i-1} \cap Z_{i-1}$. (We take M_1 to be M .) By Proposition 2.12.2, proved below, it follows that each one of the M'_i s, are rich. Observe that $G_{M_{2n-1}}$ is K_{2n-1} plus a pendent edge $[1, 2n]$: at this point vertex $2n$ is monovalent. Proposition 2.12.3, given below, shows that the deletion in M^t of a pendent edge does not alter the deficiency of the map. Therefore, N defined from M_{2n-1}^t by the deletion of $[1, 2n]$ is a rich map with $G_N = K_{2n-1}$. The proof of the theorem is thus reduced to the proofs of Propositions 2.12.1, 2.12.2, and 2.12.3. ■

The terminology involved in the statement of the following proposition is defined in the proof of Theorem 2.12.

(2.12.1) Proposition: For $i = 2, 3, \dots, 2n-1$, edge $[i, 2n]$ is an element of

a coboundary in $F_{i-1} \cap Z_{i-1}$.

Proof: Since $M_1 = M$ is rich, it follows by Lemma 2.9.3 that $F_1 \cap Z_1$ is equal to the bicycle space of K_{2n} . Consider the following sequence, B_1 , of $2n - 2$ coboundaries in the bicycle space of K_{2n} , and hence in $F_1 \cap Z_1$:

$$B_1 = (\delta(\{1, 2\}), \delta(\{1, 3\}), \dots, \delta(\{1, 2n - 1\})).$$

We also define B_j , for $2 \leq j \leq 2n - 2$, by deleting from B_1 its first $j - 1$ elements.

Observe that the only element in B_1 that contains $[2, 2n]$ is $\delta(\{1, 2\})$. The effect of deleting an edge in m^t which is not a loop in G_D nor in G_p , as in our case, is the contraction of this edge in both graphs. Therefore, the elements of $F_2 \cap Z_2$ are the elements of $F_1 \cap Z_1$ which do not contain edge $[2, 2n]$. Hence, every element in B_2 is an element of $F_2 \cap Z_2$.

The above argument iterates as follows: the only coboundary in B_j which contains edge $[j+1, 2n]$ is its first coboundary, $\delta(\{1, j+1\})$. Therefore, the elements of B_{j+1} are also elements of $F_{j+1} \cap Z_{j+1}$. The proposition follows because for $i = 2, 3, \dots, 2n - 1$ edge $[i, 2n]$ is an element of coboundary $\delta(\{1, i\})$, which is a member of $F_{i-1} \cap Z_{i-1}$. ■

(2.12.2) Proposition: Assume that an edge e of m^t belongs to an element of $F \cap Z$. Then the deletion of e does not change the deficiency of M .

Proof: Let M^t denote the t -map obtained from M by the deletion of M . We want to prove that

$$\dim[V^\perp / (F + Z)] = \dim[V_e^\perp / (F_e + Z_e)].$$

To accomplish this, it is enough to show that

$$(a) \dim(V^\perp) = 1 + \dim(V_e^\perp);$$

$$(b) \dim(F + Z) = 1 + \dim(F_e + Z_e).$$

We claim that $\{e\}$ is not a coboundary in G_M : if $\{e\}$ were a coboundary in G_M , then e would be a loop in D^t , and so $\{e\}$ would not be in a member of $F \cap Z$, contradicting the hypothesis. Hence, we have $\dim(V^\perp) = l + \dim V_e^\perp$, establishing (a).

To prove (b), note that e is a non-loop in both graphs, G_D and G_P . The difference between these graphs and G_{D_e}, G_{P_e} is that in the latter two, e is contracted. Therefore, $\dim(F) = \dim(F_e) + 1$, and $\dim(Z) = \dim(Z_e) + 1$. To conclude the proof of the proposition, it suffices to show that $\dim(F \cap Z) = \dim(F_e \cap Z_e) + 1$.

The elements of $F_e \cap Z_e$ are the coboundaries of $F \cap z$ which do not contain edge e . Take an arbitrary basis of $F \cap Z$:

$$\{A_1, \dots, A_k; B_1, \dots, B_\ell\}$$

where the A_i 's do not contain e , and the B_i 's contain e . Observe that $\ell \leq 1$. Replace this basis by

$$\{A_1, \dots, A_k; B_1, B_1 + B_2, \dots, B_1 + B_\ell\}.$$

In the new basis the only member that contains e is B_1 . Therefore, if we remove B_1 from the above set, we have a basis for $F_e \cap Z_e$ proving that $\dim(F \cap Z) = 1 + \dim(F_e \cap Z_e)$. The proof of the proposition is complete. ■

(2.12.3) Proposition: If e is a pendent edge of G_M then the deletion of e in m^t does not change $\text{def}(M)$.

Proof: Denote by M^t the t -map obtained from M^t by deleting e . Edge e is a loop in both G_D and G_P . Therefore, $F + Z$ is equal to $F_e + Z_e$, where F_e and Z_e are the coboundary spaces of D_e and P_e , dual and phial of M_e . Also, since $\{e\}$ is a coboundary in G_M , $V^\perp = V_e^\perp$, concluding the proof. ■

We observe that with the proof of the latter proposition we have com-

plete the proof of Theorem 2.12. Thus, all complete graphs are map-rich.

If we let the torus take the role of the projective plane, a parallel development shows that all complete bipartite graphs are map-rich. Also using the torus we can show that the complete tripartite graphs of the form $K_{n,n,n}$ are map-rich. Those are straightforward applications of Theorem 2.9 and the techniques developed in the case of the complete graphs.

With the help of Theorem 2.9 it is an easy matter to get examples of rich maps in arbitrary surfaces: draw a system of lines which generates the homology (mod 2) of the surface: draw enough more lines to have a 2-face-colorable t -map; deform locally some lines incident to a multiple crossing to obtain a 4-regular graph. Indeed, by the equivalence between (i) and (iii) in part (a) of theorem 2.9, every rich-map arises in this way.

Now we use the equivalence between (i) and (ii) in Theorem 2.9(a) to show that not all graphs are map-rich. For a graph G denote by G^d the graph obtained from G by doubling each one of its edges.

(2.13) Lemma: For any given connected graph G , G^d is map-rich iff it has an sp -embedding in the plane.

Proof: Observe that the coboundary space V of G^d is contained in its cycle space $V^\perp$. Therefore, $\dim(V \cap V^{perp}) = \dim(V) = v - 1$, where $v = |VG^d|$. It follows from Theorem 2.9 (ii) that there exists a rich map P with $G'_P = G^d$ iff $\chi(S_p) = (3 - v) + (v - 1)$, or $\chi(S_p) = 2$. Thus, G^d is map-rich iff it has an sp -embedding in the plane, proving the lemma. ■

Using the above lemma we can prove easily by exhaustion that if G is the Petersen graph, then G^d is not map-rich. This exhaustion is made easy by the symmetry of the Petersen graph and by the fact that C^d where C is a polygon without chords, has a unique sp -embedding in the plane.

We conclude this chapter with the establishment of an upper bound for the maximum number of independent vertices of a map-rich graph G which

has all its vertices odd valent.

(2.14) Theorem: Assume that all the vertices of a map-rich graph G , connected and without loops, are odd valent. Denote by γ the dimension of the bicycle space of G and let $v = |VG|$. The maximum number of independent vertices in G , α , is at most equal to $v - l - \gamma$. In particular, if G is planar, then $\alpha \leq v - z$, where z is the number of zigzag paths in a planar realization of G .

Proof: Take a rich map M such that $G_M = G$. As usual, denote by P the phial of M ; hence, $P' = (H_p, S_p)$ is an sp -embedding of G . By Theorem 2.10 $\xi(S_p) = v - l - \gamma$. Take an independent set of vertices in VG , denoted I , of cardinality α . To the vertices in I correspond a collection J' of disjoint smooth paths of odd length in P' . The collection J' corresponds to a collection J of disjoint odd polygons in C_p . (Think in terms of the expansion of the vertices of H_p to form the discs bounded by the squares of P). From Theorem 1.8, it follows that the mod 2 sum of the edges in an arbitrary set of disjoint odd polygons in C_p is not equal to the mod 2 sum of the edges in some alternately bicolored polygons of C_p . Thus, the members of J are representatives of distinct classes in the quotient space

$$\frac{CS(C_p)}{\langle SQ(P) \rangle + \langle VG(P) \rangle + \langle fG(P) \rangle},$$

which by Proposition 2.6 (applied to the t -map (C_p, S_p)) has $\xi(P)$ classes. Hence,

$$\alpha = |J| \leq \xi(P) = v - 1 - \gamma.$$

If G is planar, and M is a planar map with $G_M = G$ we specialize as follows. By the corollary of 2.6, M is rich and the bicycle space of G is equal to the coboundary space of G_P , where P is the phial of M . Thus $\gamma = z - 1$ and applying the bound obtained we get $\alpha \leq v - z$. The proof is complete. ■

As two examples of planar graphs where the upper bound of Theorem 2.14 holds as equality we mention the cube and the tetrahedron: the cube

has eight vertices, four zigzags (in the planar drawing) and α equal to 4; the tetrahedron has four vertices, three zigzags and α equal to 1.

Chapter 3

Gauss Code Problem Beyond the Plane

3.A Preliminary Concepts and Relation with Map-Rich Graphs

The Gauss code problem is the following: given a cyclic sequence of symbols in which each symbol occurs twice, find necessary and sufficient conditions to embed a closed curve in the plane with the restriction that the cyclic sequence of self-intersections of the curve reproduces the given cyclic sequence, when the symbols are identified with the self-intersection points. Consider, for instance, the cyclic sequence $(1, 3, 4, 6, 7, 2, 3, 9, 6, 5, 2, 1, 8, 7, 5, 4, 9, 8)$.




Figure 3.1

A planar “realization” of this sequence is given in Figure 3.1. Topological as well as combinatorial solutions for this problem are known. See a brief history in Grünbaum [10]. Combinatorial solutions are provided by Lovasz and Marx [17] and by Rosenstiehl [19]. In this chapter we consider a variant of this problem which replaces the plane for an arbitrary surface, but introduces the restriction that the t -map formed by the embedding of the curve is 2-face-colorable. For the plane the 2-face-colorability is automatically satisfied since the dual of a 4-regular planar graph is bipartite. We present a necessary and sufficient algebraic condition (Corollary of (3.3.a),(3.4), and (3.5.a)) for the realizability of a given sequence in a given surface, in which the restriction of 2-face-colorability is imposed. The conditions provide a good combinatorial algorithm for the realizability problem in the cases of surfaces with connectivity at most one: plane and projective plane.

In this section we introduce a few more concepts arising from maps and we show how the solution of the extended version of the Gauss code problem relates to map-rich graphs. A for a map M is a partition of its corners into two classes satisfying the following conditions:

- (i) For every corner x of M , x and $v_M(x)$ are in different classes.
- (ii) For every corner x of M , x and $a_M(V_M(x))$ are in the same class.

If $\{A, B\}$ is a balancing partition for map M , then the v -ordering induced by A (by B) is the collection of cyclic sequences formed by the squares containing the corners in the cycles of permutation $a_M \circ V_M$ acting on A (on B).

Observe that the elements in the v -ordering induced by A are the inverse cyclic sequences of the elements in the v -ordering induced by B .

A square of M is called relative to a given balancing partition if opposite corners of the square belong to the same class of the balancing partition; otherwise the square is called *unbalanced*.

A set of squares of M is called an *imbalance* in M if it is the set of

unbalanced squares relative to some balancing partition.

We note that if $\{A, B\}$ is a balancing partition for M , then interchanging between A and B the corners incident to a v -gon of M produces a balancing partition (A_1, B_1) . Moreover, it is evident that given two balancing partitions for M we can go from one to the other by the above interchanges. This observation implies that the set of imbalances of a map M corresponds, under the identification of $SQ(M)$ with eG_M to a coset of the coboundary space of G_M . It implies also that M is orientable iff the null set of edges is an imbalance of M ; in the latter case (and only in it) the set of imbalances in M is the coboundary space of G_M .

A *descriptor* for a map M is a v -ordering relative to some A , where $\{A, B\}$ is a balancing partition for M , together with the imbalance relative to $\{A, B\}$.

The following proposition justifies the terminology “descriptor”.

(3.0) Proposition: Every map is reconstructible from a descriptor for it.

Proof: To define a map M we simply have to specify how the corners of its squares are paired by its a_M -edges. Use the v -ordering of the descriptor to pair consistently (with the v -ordering) the corners of the squares. Clearly, there are many different ways to effect a consistent pairing. Choose an arbitrary such pairing and call the resulting map N . Choose a balancing partition for N and denote by I the induced imbalance. Denoting by J the imbalance of the given descriptor, what we have to do to transform N into M is to interchange the incidences of the pairing edges at the ends of one v_M -edge for each square in the symmetric difference of I and J . All those interchanges are necessary to satisfy the given imbalance. They are also sufficient because the descriptor constructed is a descriptor for the resulting map. ■

Edmonds’ embedding technique [4], in our language, describes an orientable map as a v -ordering and implicitly lets its imbalance be the null

set.

In the remaining of this section we show how the problem of deciding whether a given graph with maximum valency 3 is map-rich is reduced to the problem of deciding what is the minimum connectivity of a surface where a realization of a Gauss code satisfying the 2-face-colorability exists.

The material that follows in the rest of this section is not a pre-requisite for Section 3.

Given a connected graph G , the algorithmic problem of deciding whether G is map-rich seems to be difficult. A partial answer is given by Theorem 2.9: G is map-rich iff it admits an sp -embedding in the surface S of Euler characteristic $\chi(S) = 3 - |VG| + \gamma$, where γ is the dimension of the bicycle space of G , and orientability label $OL(S) = 1$ or -1 according to whether or not G is eulerian. Observe that γ can be computed by an algorithm, polynomial in the size of G : denote by V the coboundary space (over $GF(2)$) of G . Whence, $\gamma = \dim(V \cap V^\perp) = |eG| - \dim(V + V^\perp)$; the latter equality follows from 2.0.1 (c) and (d). The computation of $\dim(V + V^\perp)$ reduces to the problem of finding the rank of a matrix whose rows are formed by the characteristic vectors of arbitrary bases for V and $V^\perp$.

However, knowing the surface where G must be sp -embedded to establish that G is map-rich still leaves open the problem of finding the sp -embedding.

The situation is analogous to graphs that have an abstract dual. Given a graph G , the easiest way to establish that G has an abstract dual is, by Whitney's Theorem [26], to construct a (usual) embedding in the plane. Since there are good algorithms to test planarity, the problem of deciding whether a graph G has an abstract dual is well-solved.

Part of the problem of testing sp -embeddability of a given graph in a specific surface lies in the fact that we do not know which cyclic sequence the points of intersection of a smooth path (that in the original graph corresponds to a vertex) will define.

In view of the above we were led to consider the problem of deciding whether a given graph has an *sp*-embedding in a specific surface, with the restriction that the inverse pairs of cyclic sequences of intersection points of the smooth paths are fixed in advance. For graphs of maximum valency 3 this problem is equivalent to the unrestricted one, since these graphs have just one possibility for the inverse pairs of cyclic sequences.

Now we show how the restricted problem is algorithmically reduced to its particular case in which the graph has just one vertex, or equivalently, in which there is just one smooth path forming the *sp*-embedding. Assume that a connected graph G is *sp*-embedded in a surface S . Consider that G has more than one vertex. Take an edge e of G which is not a loop. Edge e is the intersection point of two distinct smooth paths in the *sp*-embedding. There are two ways of replacing e by two points, e' and e'' , (which become the vertices of a bounding digon in S), such as to merge the two smooth paths into one. See Figure 3.2.




Figure 3.2

Assume that R and T are subsequences of crossings points of the smooth paths that cross at e which together with e form the complete cyclic sequences of the crossings points of the same smooth paths. Combinatorially, the two topologically distinct ways to merge the smooth paths correspond to the replacement of the pair of cyclic sequences $\{(e, R), (e, T)\}$ by one of the single cyclic sequences: (e', e'', R, e', e'', T) or $(e', e'', R, e', e'', T^{-1})$, where T^{-1} represent the inverse of sequence T .

Observe that there is no possibility of real distinction between the two ways of merging the smooth paths. But since both of them work well, in the

sense that the operation does not alter the surface S , we simply choose one of the two possibilities arbitrarily. By repeating this process we reach the point in which there is just one cyclic sequence. An sp -embedding which realizes this cyclic sequence, in a minimum-connectivity surface, has the new digons introduced as bounding digons. By shrinking to a point these bounding digons we obtain an sp -embedding of the original graph.

By the above reduction and by Theorem 2.10, which gives a lower bound on the connectivity of a surface in which there exists an sp -embedding of a given graph, the problem of determining whether a graph of maximum valency 3 is map-rich would be solved if we knew an algorithm to determine what in the minimum connectivity of a surface in which an sp -embedding of a graph with one vertex and given cyclic sequence of edges around its vertices exists.

With this reduction we conclude our analysis of the connection between Gauss codes and map-rich graphs. In the next section we no longer consider map-rich graphs.

3.B Main Theorem and Crossing Functions

We introduce some formal definitions related to the Gauss code problem.

A *Gauss code* X is a cyclic sequence in which each symbol appears twice. The set of symbols of X is denoted by $\text{symb}(X)$.

A Gauss code X is *realizable* in a surface S if there exists a map P with one v -gon, such that X is equal to one of the two v -orderings of P , and such that $S = S_M$, where M is the phial of P .

Observe that if P formally realizes X in the sense of the above definition, then M' (the medial of the phial of P), is the intuitive topological realization of X , as shown in Figure 3.1.

If S is the plane, then the decision of realizability is the classical Gauss code problem. In this section we extend the solution of the realizability problem to the case of the projective plane. We answer the following more general question: is this (given) Gauss code realizable in a surface S with $\xi(S) \leq 1$? The particularization of our conditions to the plane reproduces Rosenstiehl's conditions [19]. Indeed, the methods employed generalize those of [19].

A Gauss code gives the v -ordering of a map. In order to have a complete description of a map we need to have, by Proposition 3.0, and the definition of descriptor, the associated imbalance. This indicates that to find a convenient 2-coloration of $\text{symb}(X)$ (defined next) is our aim in the (generalized) Gauss code problem.

A is a Gauss code X in which $\text{symb}(X)$ is partitioned into “black” and “white” elements. We then let the set of white elements in $\text{symb}(X)$ be the imbalance of the map whose v -ordering is given by X . Therefore, we identify a 2-colored Gauss code with a map with one v -gon.

If P is a map with one v -gon, let us denote by one of the two v -orderings of P . $\overline{P}$ is, therefore, identified with a (not yet 2-colored) Gauss code. The input for the problem is $\overline{P}$ and we seek a 2-coloration of $\text{symb}(\overline{P})$ such that the map defined P has phial M with the appropriate surface.

Observe that in the definition of realizability we can replace P by $P^\sim$. It is easy to check (we do it in Proposition 3.2.1) that the complement of each imbalance in P is an imbalance in $P^\sim$. These observations imply that realizability is independent of the order of terms of the {black,white} partition.

We also remark that any {black,white} partition of a 2-colored Gauss code P has an intuitive interpretation in M' , even though we do not use this interpretation in the sequel. In M' the elements of $\text{symb}(\overline{P})$ are vertices of H_M . Suppose that we orient the edges of H_M in the direction of a traversal of the unique smooth path of M' . Then, the 2-coloration of the faces of M'

induces a 2-coloration of the vertices of H_M : a vertex x of H_M is a black (white) vertex, whence a black (white) element in $\text{symb}(\overline{P})$, if two of the darts directed toward x or away from x encompasses a black (white) face of M' . For instance, in Figure 3.1 the black elements are $\{2, 3, 7, 9\}$ and the white elements are $\{1, 4, 5, 6, 8\}$. Thus, we see that an arbitrary 2-coloration of a Gauss code is realizable in the above sense.

Given a Gauss code X , we denote by $i_X(p)$, for $p \in \text{symb}(X)$, the subset of $\text{symb}(X)$ which is *interlaced* with p . An element $p \in \text{symb}(X)$ is said to be interlaced with $q (\neq p)$ if (p, q, p, q) is a subsequence of X . Note that since X is cyclic, the relation i_X is symmetric.

The following theorem and its corollary are the main results of the chapter.

(3.1) Theorem: A Gauss code X is realizable in a surface of connectivity at most 1 if and only if it is possible to color $\text{symb}(X)$ with two colors such as to satisfy the following condition:

(EQ) For every pair y, z of elements in $\text{symb}(X)$ the following equivalence holds:

$$|i_X(y) \cap i_X(z)| + |i_X(y)| \cdot |i_X(z)|$$

is odd iff y and z are interlaced and have the same color.

We want to point that condition (EQ) was suggested by D.H. Younger. Theorem 3.1 has a corollary which gives a good algorithm to decide if the 2-coloring exists, and in this case the corollary also provides all the 2-colorings which makes (EQ) true.

To state the corollary of Theorem 3.1 it is convenient to define the concept of interlace graph induced by a Gauss code. Given a Gauss code X the *interlace graph* induced by X , denoted by I_X , is the simple graph whose set of vertices is $\text{symb}(X)$ and whose set of edges is formed by the pairs of elements in $\text{symb}(X)$ which interlace.

Corollary of (3.1). (Good algorithm for the realizability of Gauss codes in surfaces with $\xi \leq 1$): Given a Gauss code X there exists a 2-coloration which satisfies (EQ) of Theorem 3.1 if and only if the following conditions hold:

- (i) For every pair $\{y, z\}$ of elements in $\text{symb}(X)$ which is not an edge of I_X the sum $|i_X(y) \cap i_X(z)| + |i_X(y)| \cdot |i_X(z)|$ is even.
- (ii) The subset of eI_X given by

$$T = (\{y, z\} \in eI_X : |i_X(y) \cap i_X(z)| + |i_X(y)| \cdot |i_X(z)| \text{ is even})$$

is a coboundary in the interlace graph I_X .

Moreover if (i) and (ii) hold, then the 2-colorings of $\text{symb}(X)$ which make (EQ) true are precisely the ones of the form $\{B, W\}$, where $T = \delta_{I_X}(B)$ and $W = \text{symb}(X) \setminus B$.

Proof: Initially we show that if conditions (i) and (ii) are true, then the rule to form the 2-colorings $\{B, W\}$ makes (EQ) true. Thus, let $\{B, W\}$ be a suitable 2-coloring. The pairs of equal elements always satisfy (E0), independently of the 2-coloring. If $\{y, z\}$ is not an edge of I_X , then (i) implies that $|i_X(y) \cap i_X(z)| + |i_X(y)| \cdot |i_X(z)|$ is even; hence, (EQ) is verified. If $\{y, z\} \in eI_X$, then condition (ii) and the rule to obtain $\{B, W\}$ imply that y and z are of different colors iff $\{y, z\} \in T$, i.e., $|i_X(y) \cap i_X(z)| + |i_X(y)| \cdot |i_X(z)|$ is even. Observe that also in this case (EQ) holds.

Suppose now that $\{B, W\}$ is a 2-coloring that satisfies (EQ). If $\{y, z\} \notin eI_X$, then (EQ) implies that $|i_X(y) \cap i_X(z)| + |i_X(y)| \cdot |i_X(z)|$ is even; hence, (i) is verified. By (EQ), the subset of eI_X whose elements have ends of different colors is precisely T . This implies that the graph obtained from I_X by contracting the edges not in T is bipartite. Whence, T is a coboundary in T_X , proving that (ii) holds. The proof of the corollary is complete. ■

Observe that, by the above corollary the number of essentially distinct

ways to choose the 2-coloring, and thus realize the Gauss code, is $2^{p(I_X)-1}$ where $p(I_X)$ is the number of components of I_X .

By a result that we prove later in the chapter, (Corollary of 3.3(a), 3.4, and 3.5(a), presented immediately after the proof of Lemma 3.5,) the surface where a Gauss code X is realizable is orientable iff I_X is eulerian. This property permits the following specialization for the plane of the conditions of the above corollary:

Rosdnstiehl's conditions: A Gauss code X is realizable in the plane if and only if the following conditions hold:

- (I) I_X is eulerian;
- (II) for $\{y, z\} \notin eI_X$, $|i_X(y) \cap i_X(z)|$ is even;
- (III) $T = \{\{y, z\} \in eI_X : |i_X(y) \cap i_X(z)| \text{ is even}\}$ is a coboundary in I_X .

Conditions (I), (II), (III) appear in Rosenstiehl [19]. The proof of Theorem 3.1 takes most of the remaining of the chapter. To establish the proof we develop a theory of “crossing functions” which generalizes for arbitrary surfaces the “projections” described for the plane by Shank in [24].

Given a 2-colored Gauss code X we now define the *crossing function* induced by X

$$c_X : 2^{\text{symb}(X)} \longrightarrow 2^{\text{symb}(X)}.$$

Function c_X is given on the singletons by $c_X(\{a\}) = i_X(a) + t_X(a)$, where

$$t_X(a) = \begin{cases} \phi & \text{if } a \text{ is white} \\ \{a\} & \text{if } a \text{ is black,} \end{cases}$$

and extended for arbitrary $A \subseteq \text{symb}(X)$ as follows: $c_X(A) = \sum_{a \in A} c_X(\{a\})$, where the sum is mod 2 sum of subsets. It is also convenient to extend t_X and i_X to $2^{\text{symb}(X)}$ by the analogous formulas: for $A \subseteq \text{symb}(X)$,

$$t_X(A) = \sum_{a \in A} t_X(a)$$

$$i_X(A) = \sum_{a \in A} i_X(a)$$


In the proof of Theorem 3.1 we denote the abstract Gauss code X by $\overline{P}$, with the intention of working with the terminology of maps that we have developed.

Given a Gauss code $\overline{P}$, denote by $ODD(\overline{P})$ and $EVEN(\overline{P})$ the following partition of $\text{symb}(\overline{P})$:

$$ODD(\overline{P}) = \{y \in \text{symb}(\overline{P}) : |i_P(y)| \text{ is odd}\}$$

$$EVEN(\overline{P}) = \{y \in \text{symb}(\overline{P}) : |i_P(y)| \text{ is even}\}$$

To simplify the notation, we use $i_P(X), c_P(X), t_P(X)$, instead of the more correct $i_{\overline{P}}(\{x\}), c_P(\{x\}), t_{\overline{P}}(x)$.



Dependence Structure for Theorem 3.1

Proof of Theorem 3.1. Lemma 3.2, proved below, states that a Gauss code $\overline{P}$ is realizable in S with $\xi(S) \leq 1$ iff there exists a 2-coloring of $\text{symb}(\overline{P})$ inducing a map P with one v -gon, (the 2-coloring gives the imbalance relative to the v -ordering given by $\overline{P}$), and a crossing function c_P which satisfies the following equations:

$$(a) \quad (c_P + c_P^2)(x) = \phi, \quad \text{for } x \in \text{EVEN}(\overline{P});$$

$$(b) \quad (c_P + c_P^2)(x) = \text{ODD}(\overline{P}), \quad \text{for } x \in \text{ODD}(\overline{P}).$$

We assume Lemma 3.2 and show the equivalence of condition (EQ) of Theorem 3.1 with conditions (a) and (b) of Lemma 3.2. We begin by deriving a useful expression for $(c_P + c_P^2)(x)$. For $z \in \text{symb}(\overline{P})$ let the variable $\gamma_z \in \{0, 1\} = GF(2)$ assume value 1 if z is black and 0 if z is white. Since $c_P = i_P + t_P$ and $t_P^2 = t_P$, we obtain

$$\begin{aligned} (c_P + c_P^2)(x) &= i_P^2(x) + i_P t_P(x) + t_P i_P(x) + i_P(x) \\ &= i_P^2(x) + \sum_{y \in i_P(x)} (\gamma_x + \gamma_y + 1)y. \end{aligned}$$

It is convenient to separate the equivalence (EQ) into two equivalences (a') and (b') stated below:

(a') if $|i_P(x)| \cdot |i_P(z)|$ is even, then $|i_P(x) \cap i_P(z)|$ is odd iff $\gamma_x = \gamma_z$ and $x \in i_P(z)$;

(b') if $|i_P(x)| \cdot |i_P(z)|$ is odd, then $|i_P(x) \cap i_P(z)|$ is even if $\gamma_x = \gamma_z$ and $x \in i_P(z)$.

It is clear that statements (a') and (b') together are equivalent to (EQ) of Theorem 3.1. Our aim in the remaining of the present proof is to establish the equivalence

$$(a) \quad \text{and} \quad (b) \Leftrightarrow (a') \quad \text{and} \quad (b').$$

A helpful remark in this direction is that for $A \subseteq \text{symb}(\overline{P})$, $z \in i_P(A)$ iff $|i_P(z) \cap A|$ is odd. Whence, $z \in i_P^2(x)$ iff $|i_P(z) \cap i_P(x)|$ is odd.

With the above remark and the symmetry of i_P and of i_P^2 , (a') can be restated as follows: if $x \in EVEN(\overline{P})$, then $z \in i_P^2(x)$ iff $\gamma_x = \gamma_z$ and $x \in i_P(z)$. This statement, by the expression calculated above for $(c_P + c_P^2)(x)$, is equivalent to the following one: if $x \in EVEN(\overline{P})$, then $(c_P + c_P^2)(x) = \phi$, which is precisely statement (a) . Therefore, we have established the equivalence $(a) \Leftrightarrow (a')$.

To conclude, we show the equivalence

$$(a) \quad \text{and} \quad (b) \Leftrightarrow (a) \quad \text{and} \quad (b').$$

Statement (b') can be rewritten as follows: if x and z are both in $ODD(\overline{P})$, then $z \in i_P^2(x)$ iff $\gamma_x = \gamma_z$ and $x \in i_P(z)$. By the expression for $(c_P + c_P^2)(x)$, this shows that (b') is equivalent to the following statement: if x and z are both in $ODD(\overline{P})$, then $z \in (c_P + c_P)(x)$. It is then clear that (b) implies (b') and that (a) , the symmetry of the relation $c_P + c_P^2$, and (b') imply (b) .

This concludes the Proof of Theorem 3.1, provided Lemma 3.2 is established. ■

(3.2) Lemma: A Gauss code $\overline{P}$ is realizable in surface S of connectivity at most one if and only if there exists a 2-coloring of $\text{symb}(\overline{P})$ inducing a map P with a crossing function c_P which satisfies the following equations:

$$\begin{aligned} (a) \quad & (c_P + c_P^2)(x) = \phi, \quad \text{for } x \in EVEN(\overline{P}); \\ (b) \quad & (c_P + c_P^2)(x) = ODD(\overline{P}), \quad \text{for } x \in ODD(\overline{P}); \end{aligned}$$

Proof: Assume that there exists a 2-coloring of $\text{symb}(\overline{P})$ such that equations (a) and (b) hold. These equations imply that the dimension of the image of $c_P + c_P^2$, $\dim[Im(c_P + c_P^2)]$, is at most one. The corollary of Proposition 3.2.1, which we prove next, states that $c_P + c_{P\sim}$ is the identity. Whence, $c_{P\sim} \circ c_P = c_P + c_P^2$ and thus, $\dim[Im(c_{P\sim} \circ c_P)] \leq 1$. By Theorem 3.3(a), proved below, we have $\xi(M) = \dim[Im(c_{P\sim} \circ c_P)]$, where M is the phial of P . Therefore, $\xi(M) \leq 1$, and this proves that P is realizable in a surface of

connectivity at most one. This part of the proof is complete, provided the corollary of Proposition 3.2.1 as well as Theorem 3.3(a), are established.

Conversely, assume that the Gauss code $\overline{P}$ is realizable in the plane or in the projective plane. This means that there exists a 2-coloring of $\text{symb}(\overline{P})$ such that the map P formed has phial M satisfying $\xi(M) \leq 1$. For such maps, Proposition 3.2.2, which we prove below, states that the s -cycles in $M^\perp$ are precisely the elements of the coboundary space F of G_D (where D is the dual of M). Lemma 3.4, also proved below, states that for maps M whose phial P has one v -gon $c_P(x)$ is an s -cycle in M^t if and only if $|i_P(x)|$ is even, i.e., $x \in \text{EVEN}(\overline{P})$. Putting together 3.2.2 and 3.4 we obtain the following condition for maps P with one v -gon and $\xi(M) \leq 1$: $c_P(x)$ is in F iff $x \in \text{EVEN}(\overline{P})$.

Lemma 3.5(b), proved below, when applied to map P states that $\text{Ker}(c_{P^\sim}) = F$. Therefore, $c_P(x)$ is in $\text{Ker}(c_{P^\sim})$ iff $x \in \text{EVEN}(\overline{P})$. This condition implies $(c_{P^\sim} \circ c_P)(x) = \phi$, for $x \in \text{EVEN}(\overline{P})$. Since, as mentioned before $c_{P^\sim} \circ c_P = c_P + c_P^2$, this concludes (a), modulo the unproved statements, whose proofs are supplied in the sequel.

To prove (b) consider $x \in \text{ODD}(\overline{P})$. Recall that

$$(c_P + c_P^2)(x) = i_P^2(x) + \sum_{y \in i_P(x)} (\gamma_x + \gamma_y + 1)y,$$

an expression that we computed in the Proof of Theorem 3.1. Also recall that for $z \in \text{symb}(\overline{P})$, $\gamma_z = 1$ iff z is a black element of $\text{symb}(\overline{P})$. Therefore, $y \in (c_P + c_P^2)(x)$ iff one of the three following conditions holds:

- (i) $y \in i_P^2(x)$, $y \in i_P(x)$ and $\gamma_x \neq \gamma_y$;
- (ii) $y \in i_P^2(x)$, and $y \notin i_P(x)$;
- (iii) $y \notin i_P^2(x)$, $y \in i_P(x)$ and $\gamma_x = \gamma_y$.

If $y \in \text{EVEN}(\overline{P})$ we show that $y \notin (c_P + c_P^2)(x)$. Note that $y \in (c_P \circ c_P^2)(x)$ iff $x \in (c_P + c_P^2)(y)$, property easily deduced from (i), (ii), and

(iii) above. But $(c_P + c_P^2)(y)$ is the null subset, by part (a). Thus, if $y \in \text{EVEN}(\overline{P})$, then $y \notin (c_P + c_P^2)(x)$. To establish (b) it suffices, then, to prove that $y \in (c_P + c_P^2)(x)$, for $y \in \text{ODD}(\overline{P})$. To this end initially observe that $z \in (c_P + c_P^2)(z)$, for every $z \in \text{ODD}(\overline{P})$. This follows from $z \in i_P^2(z)$ and $z \notin i_P(z)$. Since $\xi(M) \leq 1$ and $c_P + c_P^2 = c_{P^\sim} \circ c_P$, Lemma 3.3(a) proved below implies that $\dim[\text{Im}(c_P + c_P^2)] \leq 1$. Whence, there are one or two elements in $\text{Im}(c_P + c_P^2)$ one of which is the null vector. Thus, we conclude that for $y \in \text{ODD}(\overline{P})$, $(c_P + c_P^2)(x) = (c_P + c_P^2)(y)$, because both images are non-null. Now to finish the proof assume an arbitrary $y \in \text{ODD}(\overline{P})$. We show that at least one of the conditions (i), (ii), (iii) above is verified. Suppose, as a first case that $y \in i_P^2(x)$. If $y \notin i_P(x)$, then (ii) is verified. Therefore, assume that $y \in i_P(x)$. It $\gamma_x = \gamma_y$, then (i) is verified. Suppose then, that $\gamma_x \neq \gamma_y$. We are under the hypotheses $y \in i_P^2(x)$, $y \in i_P(x)$ and $\gamma_x \neq \gamma_y$. By the expression for $(c_P + c_P^2)(x)$ this implies that $y \notin (c_P + c_P^2)(x)$. But we know that $y \in (c_P + c_P^2)(y)$ and that $(c_P + c_P^2)(x) = (c_P + c_P^2)(y)$. Therefore, $y \in (c_P + c_P^2)(x)$, which is a contradiction, concluding the analysis in the case that $y \in i_P^2(x)$. Assume the second case, $y \notin i_P^2(x)$. Since $y \in (c_P + c_P^2)(x)$ we conclude that $y \in i_P(x)$ and that $\gamma_x \neq \gamma_y$. This is precisely condition (iii). This concludes the proof of (b). The proof of the lemma is, then, reduced to the proofs of the Corollary of 3.2.1 and of the following statements: 3.2.2, 3.3(a), 3.4, and 3.5(b). ■

For the next proposition and its corollary we consider, as usual, $SQ(X) \cong SQ(M)$ if $X \in \Gamma(M)$.

(3.2.1) Proposition: Given any map M , $T \subseteq SQ(M)$ is an imbalance in M iff $SQ(M) \setminus T$ is an imbalance in $M^\sim$.

Proof: Assume that T is an imbalance in M . By definition there exists a balancing partition B_T for M which has T as its imbalance. Since $M^\sim$ has the same v -gons as M , B_T is also a balancing partition for $M^\sim$. Given a corner X of M or of $M^\sim$, X and $v_M(X) = v_{M^\sim}(X)$ are not in the same class of B_T . This implies that two corners of a square which are in the same class

of B_T are adjacent in C_M iff they are opposite in the corresponding square of $M^\sim$. The converse follows by the same argument, since the operations of taking the antimap of a map and taking the complement of a subset are involutions. ■

Corollary of Proposition 3.2.1: For a map P with one v -gon, the linear transformation

$$c_P + c_{P^\sim} : 2^{SQ(P)} \longrightarrow 2^{SQ(P)}$$

is the identity.

Proof: The corollary follows from the definitions of c_P and $c_{P^\sim}$, because the above proposition implies that a symbol of the associated 2-colored Gauss codes is black in P iff it is white in $P^\sim$. ■

(3.2.2) Proposition: Given a map M , if $\xi(M) \leq 1$, then the s -cycles of M^t are precisely the elements of the coboundary space F of G_D , where D is the dual of M .

Proof: If $\xi(M) = 0$, the proposition follows from (2.6), since the hypothesis implies that $V^\perp = F$. If $\xi(M) = 1$, then by Theorem 1.7 M is non-orientable, since the parity of $\xi(M)$ and $\chi(M)$ agree. Take an odd polygon in C_M and consider its image under ψ_c^M . This image is an r -cycle in M^t , which we denote by $4x$. The cycle $4x$ is not in F , since the elements of F are s -cycles. Therefore, by Proposition 2.6, it follows that

$$V^\perp/F = \{F, x + F\}.$$

Consider an s -cycle y . Either $y \in F$ or else $y \in x + F$. In the latter case, $x + y \in F$. Since ψ_c^M is a homomorphism, by Proposition 1.4, the symmetric difference of an r -cycle and an s -cycle is an r -cycle. Hence, $x + y$ is an r -cycle which is contained in F . This is not possible. Therefore, $y \in F$, establishing the proposition. ■

3.C Properties of Crossing Functions

In this section we continue the proof of the main theorem (Theorem 3.1). At the same time we establish some theorems about general crossing functions associated with t -maps with one vertex. Items (b) and (c) of the next theorem are not used in connection with Theorem 3.1, but are included for completeness.

(3.3) Theorem: if P is a map with one v -gon, then we have

- (a) $\dim[Im(c_{P^\sim} \circ c_P)] = \xi(M);$
- (b) $Im(c_{P^\sim} \circ c_P) = V^\perp \cap F^\perp;$
- (c) $Ker(c_{P^\sim} \circ c_P) = V + F,$

where M is the phial of P , D is the dual of M , and V and F are the coboundary spaces of G_M and G_D , respectively.

Proof: To prove this theorem we make use of Lemma 3.5, proved below. By Lemma 3.5(a) we have $Im(c_P) = V^\perp$. Therefore,

$$Im(c_{P^\sim} \circ c_P) = Im(c_{P^\sim}|V^\perp),$$

where, “ $|V^\perp$ ” stands for restriction to $V^\perp$. By the fundamental theorem for homomorphisms applied to $c_{P^\sim}|V^\perp$ we have

$$\dim[Im(c_{P^\sim}|V^\perp)] + \dim[rmKer(c_{P^\sim}|V^\perp)] = \dim V^\perp.$$

By lemma 3.5(b), applied to map $P^\sim$, it follows that $Ker(c_P) = F$. By Proposition 2.2, $F \subseteq V^\perp$, implying $Ker(c_{P^\sim}|V^\perp) = F$. Therefore,

$$\dim[Ker(c_{P^\sim}|V^\perp)] = f - 1,$$

where $f = |VG_D|$ (Observe that since P has one v -gon, D is connected). Let

$v = |VG_M|$ and $e = |eG_M|$. By the above equations, it follows that

$$\begin{aligned} \dim[Im(c_{P^\sim} \circ c_P)] &= (e - v + 1) - (f - 1) \\ &= (e + 2) - (v + f) \\ &= \xi(M) \end{aligned}$$

The proof of part (a) is complete, provided Lemma 3.5 is established.

We proceed to prove (b). It follows, from the Corollary of Proposition 3.2.1, that $c_{P^\sim} \circ P = c_{P^\circ}$. This commutativity and the facts that, by Lemma 3.5(b), $Im(c_P) = V^\perp$ and $Im(c_{P^\sim}) = F^\perp$ imply the inclusion $Im(c_{P^\perp} \circ c_P) \subseteq V^\perp \cap F^\perp$. By the triple inclusion property, lemma 2.9.1,

$$\dim(V^\perp \cap F^\perp) - \dim(V \cap F) = \xi(M).$$

By the absorption property, Lemma 2.5, it follows that $V \cap F = V \cap F \cap Z$, with Z coboundary space of G_P . Since $|VG_P| = 1$, Z is the null space. Therefore,

$$\dim(V^\perp \cap F^\perp) = \xi(M).$$

By part (a) $\dim[Im(c_{P^\sim} \circ c_P)] = \xi(M)$. Thus, $Im(c_P \circ c_{P^\sim})$ is a subspace of $V^\perp \cap F^\perp$ which has the same dimension as the whole space. Therefore,

$$Im(c_{P^\sim} \circ c_P) = V^\perp \cap F^\perp;$$

this concludes (b).

To prove (c) we also use Lemma 3.5. Take $x \in V + F$; say that $x = u + w$ with $u \in V$ and $w \in F$. We have

$$(c_{P^\sim} \circ c_P)(u + w) = c_{P^\sim}(c_P(u) + c_P(w)) = c_{P^\sim}(c_P(w)).$$

The latter equality follows because $c_P(u) = \phi$ by Lemma 3.5(b). As c_P and $c_{P^\sim}$ commute we have

$$c_{P^\sim}c_P(w) = c_P \circ c_{P^\sim}(w) = c_P(\phi) = \phi.$$

That $c_{P^\sim}(w)$ follows from Lemma 3.5(b) applied to map $P^\sim$. We conclude then that $V + F \subseteq \text{Ker}(c_{P^\sim} \circ c_P)$.

The value of $\dim[\text{Ker}(c_{P^\sim} \circ c_P)]$ is $e - \xi(M)$, where $e = |SQ(M)|$. This follows from the fundamental theorem for homomorphisms and from part (a). The value of $\dim(V + F) = (v - 1) + (f - 1)$, where $v = |VG_M|$, $f = |VG_D|$. This follows from 2.0.1(a), and from the fact that $V \cap P$ is the null space. Recall that $V \cap F = V \cap F \cap Z$, and Z is the null space, since Z is the coboundary space of G_P which has only one vertex. We can rewrite $(v - 1) + (f - 1)$ as $e - \xi(M)$, by the definition of $\xi(M)$. Since $V + F$ is a subspace of $\text{Ker}(c_{P^\sim} \circ c_P)$ and has the same dimension, it follows that it is equal to $\text{Ker}(c_{P^\sim} \circ c_P)$. This concludes (c) and the proof of the Theorem 3.3, provided Lemma 3.5 is established. ■

(3.4) Lemma: For every map M whose phial P has one v -gon, $c_P(x)$ is a cycle in G_M , for every $x \in \text{symb}(\overline{P})$ and

$$c_P(x) \text{ is an } s\text{-cycle in } M^t \text{ iff } |i_P(x)| \text{ is even.}$$

Proof: To prove this lemma, which is of fundamental importance in the chapter, we define a function c'_P on the corners of M with image in the set of cycles of C_M . For a corner X of M , the cycle $c'_P(X)$ is defined by the edges of C_M occurring once in the reentrant path which starts at $z_M(X)$ and proceeds by using a_{M^-} , v_{M^-} , and f_{M^-} -edges (in this order) until it reaches another corner of x , which denotes the square to which x belongs. This corner can be $v_M(X)$, in which case we close the path by using the f_M -edge which links $v_M(X)$ to $z_M(X)$; it also can be $f_M(X)$, in which case we close the path by using the v_M -edge which links $f_m(X)$ to $z_m(X)$. The fact that x is a loop in G_P and a parity argument show that the first corner of x reached by the path is not X , and is not $z_M(X)$. This completes the definition of c'_P .

We claim that the image of $c'_P(X)$ under ψ_c^M is $c_P(x)$, independently

of the corner X of x . To prove this claim observe that there is just one edge of square x in $c'_P(X)$, for X corner of x . If x is an s -loop in p^t (x is a black element of $\text{symb}(\overline{P})$), it follows from the definitions that the edge of x present in $c'_P(X)$ is an f_M -edge; otherwise, if x is an r -loop in p^t , this edge is a v_m -edge. Since the f_M -edges are not contracted under ψ (the canonical contraction that transforms (C_M, S_M) into (G_M, S_M)), square x is an edge in $c_P(x)$, in the case that x is an s -loop in p^t . Since the v_M -edges are contracted under ψ , square x is not an edge in $c_P(x)$, in the case that x is an r -loop in p^t . The other edges belonging to the image of $c'_P(X)$ under ψ_c^M correspond to the squares whose intersection with $c'_P(X)$ is a path of length 2, that is, the squares which are met once by the reentrant path defining $c'_P(X)$. Those squares are precisely the squares in $i_P(x)$. The claim that $\psi_c^M(c'_P(X)) = c_P(x)$ for every corner X of square x is, therefore, proved.

By the definition of s - and r -cycles, to complete the proof of the lemma it is enough to show that the congruence

$$|c'_P(X)| = |i_P(x)| \pmod{2}$$

holds for every square x with X corner of x .

Since C_M is cubic, the cycle $c'_P(X)$ induces a subgraph of C_M which consists of a certain number of disjoint polygons, whose set is denoted by Ω . We count the vertices of these polygons. In square x there are two corners which are vertices of a polygon in Ω . If a square is met twice by the reentrant path which defines $c'_P(X)$, then all its four corners are vertices of polygons in Ω . If a square is met once by the reentrant path, then three of its corners are vertices of a polygon in Ω . Evidently, if a square is not met by the reentrant path, none of its corners is in a polygon in Ω . Hence, the parity of the number of vertices of the polygons in Ω , which is the same as the parity of $|c'_P(X)|$, is equal to the parity of $|i_P(x)|$. This establishes the Lemma. ■

(3.5) Lemma: Let $P = P_M$ be a map with one v -gon, and V the coboundary space of G_M . We then have the following equalities for the image and the

kernel of the crossing function $c_P : 2^{SQ(P)} + 2^{SQ(P)}$ induced by P :

$$(a) \quad Im(c_P) = V^\perp$$

$$(b) \quad Ker(c_P) = V$$

Proof: By Lemma 3.4 above, $c_P(x)$ is in $V^\perp$ for every $x \in \text{ymb}(P)$. Since the image of c_P is generated by the members of

$$\{c_P(x) : x \in \text{ymb}(\overline{P})\},$$

it follows that $Im(c_P) \subseteq V^\perp$.

The proof of the converse inclusion needs some additional preparation. We give it in the next section, Lemma 3.6. We assume that (a) holds for the proof of (b).

Take $s \in \text{ymb}(\overline{P})$ and $T \subseteq \text{ymb}(\overline{P})$. We show that the scalar product $\langle \{s\}, c_P(T) \rangle$ is equal to $\langle c_P(\{s\}), T \rangle$ as follows:

$$\begin{aligned} \langle \{s\}, c_P(T) \rangle &= \sum_{x \in T} \langle \{s\}, i_P(x) \rangle + \sum_{z \in T} \langle \{s\}, t_P(x) \rangle \\ &= \sum_{x \in T} \langle i_P(s), x \rangle + \sum_{z \in T} \langle t_P(s), x \rangle \\ &= \langle c_P(s), T \rangle. \end{aligned}$$

Let $T \subseteq \text{ymb}(\overline{P})$ be an element of $Ker(c_P)$. Since $\langle x, c_P(T) \rangle = \langle c_P(x), T \rangle$ and $c_P(T) = \phi$, it follows that $\langle c_P(x), T \rangle = 0$ for every $x \in \text{ymb}(\overline{P})$. As we are assuming (a), $V^\perp$ is generated by the members of $\{c_P(x) : x \in \text{ymb}(\overline{P})\}$. It follows that T belongs to the space perpendicular to $V^\perp$ namely, V . Hence $Ker(c_P) \subseteq V$.

We first learned how to prove that $V \subseteq Ker(c_P)$ from H. Shank and subsequently we discover the following shorter proof.

Take a coboundary $T \subseteq \text{ymb}(\overline{P}) \cong e(G_M)$; thus T belongs to V . Since by part (a), for $s \in \text{ymb}(\overline{P})$, $c_P(s)$ is a cycle in G_M , it follows that

$\langle c_P(s), T \rangle = 0$ for every $s \in eG_M$. This is the same as $\langle s, c_P(T) \rangle = 0$ for every $s \in eG_M$, by the equality established above. Hence, $c_P(T) = \phi$, proving that $T \in \text{Ker}(c_P)$. This concludes the proof of the lemma, provided the inclusion $V^\perp \subseteq \text{Im}(c_P)$ is established. We treat this inclusion in Lemma 3.6. ■

The following corollary is an algebraic-combinatorial characterization of realizability of a Gauss code in a given arbitrary surface:

Corollary of (3.3.a),(3.4) and (3.5.a): A Gauss code is realizable in a surface of connectivity ξ iff there exists a {black,white} partition of $\text{symb}(\overline{P})$ such that the induced map with one v -gon P satisfies

$$\dim[\text{Im}(c_{P^\sim} \circ c_P)] = \xi.$$

Moreover, the surface where $\overline{P}$ is realizable is orientable iff $I_{\overline{P}}$ is eulerian.

Proof: The first part of the corollary, treating the connectivity, is a straightforward application of the definition of realizability and of the Theorem 3.3(a).

To establish the second part, observe that for every P induced by a {black, white} partition Lemma 3.5(a) states that the members of $\{c_P(x) : x \in \text{symb}(\overline{P})\}$ generate the cycle space $V^\perp$ of G_M where M is the phial of P . By Lemma 3.4.2, $c_P(x)$ is an s -cycle in M^t iff $x \in \text{EVEN}(\overline{P})$. Note that the mod 2 sum of cycles in G_M is an s -cycle iff the number of r -cycles is even. (Recall that $\psi_c^M : CS(M) \rightarrow CS(G_M) = V^\perp$ is a homomorphism, result proved in Proposition 1.4.) Hence, $V^\perp$ has only s -cycles iff $x \in \text{EVEN}(\overline{P})$ for every $x \in \text{symb}(\overline{P})$. But $V^\perp$ has only s -cycles iff C_M is bipartite, or iff S_M is orientable. Since S_M is the surface where $\overline{P}$ is realized (once the {black,white} partition is chosen), the corollary follows. ■

3.D Surjectivity of the Crossing Functions

In this section we conclude the proofs of the statements which are still open with the proof that $V^\perp \subseteq Im(c_P)$. Initially we define some concepts, leading to the idea of “reflexivity”.

Consider a graph L and an arbitrary, but fixed, orientation for its edges. Let π be a cyclic path in L such that each edge of L appears twice in π . Given a pair (L, π) under these conditions, we define cyclic sequence π' as follows. The symbols in π' are labels for the edges of G . The order in which they occur is given by cyclic path π . Moreover each symbol x in π' is signed “+” or “−” according to whether or not edge labelled x is traversed in the right direction (according to the fixed orientation for the edges of L).

We identify π' with a 2-colored Gauss code, where the black elements are the symbols whose two occurrences in π' are with the same sign and the white elements are the others. Functions i_π , and c_π , are, therefore, defined.

Given an abstract cyclic sequence π' in which each symbol is signed and occurs twice, we define a digraph $\Omega(\pi')$ as follows. The darts of $\Omega(\pi')$ are the symbols which occur in π' . To obtain the vertices of $\Omega(\pi')$ we start with a head and a tail for each dart and effect the following identifications, one for each successive pair of signed symbols in π' :

- if $(+x, +y)$ appears in π' , then identify head (x) and tail (y) ;
- if $(+x, -y)$ appears in π' , then identify head (x) and head (y) ;
- if $(-x, +y)$ appears in π' , then identify tail (x) and tail (y) ;
- if $(-x, -y)$ appears in π' , then identify tail (x) and head (y) .

Given (L, π) we say that cyclic path π is *reflexive* in digraph L if $\Omega(\pi') = L$.

Observe that for any pair (L, π) , $\Omega(\pi')$ can be transformed into L by identifying some of its vertices. π is reflexive in L if no identification is

necessary.

Given (L, π) the crossing function c_π , has its image contained in the cycle space of L , $CS(L)$. (To get $CS(L)$ the orientation of the edges of L plays no role.) To show this inclusion, observe that the edges between the two occurrences of an edge e in π correspond to a path π_e which is reentrant if e is traversed in opposite directions in π ; π_e is made reentrant by the composition with the path $(\text{tail}(e), e, \text{head}(e))$, if e is traversed in the same direction in π .

(3.6) Lemma: If P is a map with one v -gon, then $\{c_P(x) : x \in \text{symb}(\overline{P})\}$ generates $V^\perp$.

Proof: Consider the pair (G_M, π_z) where π_z is the unique zigzag path in M^t . Consider also a fixed orientation of the edges of G_M . Let π'_z be induced by (G_M, π_z) . We claim that $c_P = c_{\pi'_z}$, when $SQ(P)$ is identified with eG_M . The cyclic sequence of squares met by the v -gon of P and the cyclic sequence of the edges of G_M given by π'_z are equal. Thus, to prove the claim, we have only to show that an edge x of G_M is traversed in the same direction twice in π_z iff the corresponding square x is not in the imbalance of P , i.e., if edge x is an s -loop in p^t . But x is an s -loop in P^t iff for X arbitrary corner of x , $(X, z_M(X))$ and $(v_M(X), f_M(X))$ are two ordered pairs of consecutive vertices in the v -gon of P traversed in one of the two directions. Observe that $(x, z_M(X))$ and $(v_M(X), f_M(X))$ are such pairs iff edge x of G_M is traversed in the same direction twice by the zigzag. Therefore, the claim is proved.

By Proposition 3.6.1, proved below, if π_z is reflexive in G_M , then the proposition follows. Therefore, assuming 3.6.1, to conclude the proof it is enough to show that for arbitrary M^t with just one zigzag path, π_z is reflexive in G_M .

If π_z is not reflexive in G_M , then, by definition, $\Omega(\pi'_z) \neq G_M$. This means that $\Omega(\pi'_z)$ has more vertices than G_M , and to obtain G_M from $\Omega(\pi'_z)$ some identifications of vertices of $\Omega(\pi'_z)$ are necessary. It follows that there are two darts, x and y , which satisfy $\text{head}(x)\text{tail}(y)$ in G_M but not in $\Omega(\pi'_z)$.

We can choose x and y verifying this condition and, moreover, the condition that x and y are consecutive edges in the cyclic sequence of edges around the vertex $\text{head}(x) = \text{tail}(y)$ in M^t . However, note that such consecutive edges correspond to a pair of consecutive edges in π_z . Whence, by definition of $\text{head}(x) = \text{tail}(y)$, which contradicts the assumption. Thus, we conclude that π_z is reflexive in G_M . The proof of the lemma is reduced to the proof of Proposition 3.6.1, which we give next. ■

The main idea in the proof the the following proposition was obtained from Read and Rosenstiehl [21].

(3.6.1) Proposition: If π is a cyclic path in a digraph L , such that π uses each edge of L twice, then $c_{\pi'} : 2^{eL} \rightarrow CS(L)$ is an epimorphism if and only if π is reflexive in L .

Proof: To prove the proposition it is enough to show that the image of c_{π} , is the cycle space of $\Omega(\pi')$. If this is proved, it follows that if π is not reflexive in L , then c_{π} , is not an epimorphism; also conversely, if π is reflexive in L , then c_{π} , is an epimorphism.

We construct $Q(\pi)'$ from π' as follows. We start by drawing an arbitrary dart representing an occurrence of a signed symbol in π' . We proceed drawing darts in the order that they appear, without lifting the pencil from the paper, as long as this is possible. Each time that an edge occurs for the first time we must draw it as a pendent edge (the final vertex must be a new one). The rule not to lift the pencil is not possible to obey when an edge e occurring for the second time is not incident to the last vertex reached. At each such occurrence, we make a copy of the graph drawn so far, denoting it by L_e . Next, we make the necessary identification of the two vertices and proceed. At the end we have a sequence of partial graphs

$$L_{e_1}, L_{e_2}, \dots, L_{e_k}, \Omega(\pi'),$$

where edge e_i forced the i -th identification. Suppose we have k identifications.

If no identifications were made ($k = 0$), then we would have $e + 1$ vertices, where e is the number of edges of $\Omega(\pi')$. Each identification reduces the number of vertices by one. Therefore, if v is the number of vertices of $\Omega(\pi')$ we have

$$e + 1 - k = v \quad \text{or} \quad k - e + 1 = v.$$

Observe that this value of k is the dimension of $CS(L)$, since L is connected.

For every $i = 1, 2, \dots, k$ the two facts that follow are evident:

- (a) for $j < i$ $c_\pi, (e_j)$ is a cycle in L_i ;
- (b) $c_\pi, (e_i)$ is not a cycle in L_i .

Fact (a) follows from the observation that a cycle remains a cycle when identifications of vertices are performed. Fact (b) follows from the equivalence: the set of edges occurring an odd number of times in a path π is a cycle iff π is reentrant.

From (a) and (b) follow that each $c_\pi, (e_i)$ is not expressible as mod 2 sum of a subset of $\{c_\pi, (e_j) : j < i\}$. Hence,

$$c_\pi, (e_1), \dots, c_\pi, (e_k)$$

are independent cycles in $\Omega(\pi')$. Since k is the dimension of the cycle space of $\Omega(\pi')$, the $c_\pi, (e_i)$ is form a basis for this space. The proof is complete. ■

Observe that with the proof of this proposition every statement made so far in Chapter 3 is established.

We conclude the chapter with an example of a class of arbitrarily large Gauss codes which are realizable only in surfaces of connectivity at most two.

Denote by $\overline{P}_n$ the Gauss code

$$(1, 2, \dots, n, 1, 2, \dots, n).$$

(3.7) Proposition: if n is odd there is one realization of $\overline{P}_n$ in the plane, and every other realization of $\overline{P}_n$ is in the torus. If n is even there is one realization of $\overline{P}_n$ in the projective plane, and every other realization of $\overline{P}_n$ is in the Klein bottle.

Proof: Suppose we are given an arbitrary {black,white} partition, $\{B, W\}$, of $\text{symb}(\overline{P}_n)$. Let us denote by M_n the phial of the map P_n induced by the 2-colored Gauss code.

Suppose that n is odd. In this case $i_{P_n}^2(x)$ is the whole of $\text{symb}(\overline{P}_n)$. (Recall that $y \in i_{P_n}^2(x)$ and $i_{P_n}(y)$ have an odd number of common elements.) As we have

$$(c_{P_n} \circ c_{P_n}) = i_{P_n}^2(x) + \sum_{y \in i_{P_n}(x)} (1 + \gamma_x + \gamma_y)y,$$

where $\gamma = 1$ iff $z \in B$, the following implications hold:

- (i) if $x \in B$, then $(c_{P_n} \circ c_{P_n})(x) = W$;
- (ii) if $x \in W$ then $(c_{P_n} \circ c_{P_n})(x) = B$;

Therefore, except in the case where one of W and B is empty, we have $\dim[\text{Im}(c_{P_n} \circ c_{P_n})] = 2$. Since $\text{ODD}(\overline{P})$ is empty, there are no r -circuits in G_{M_n} , by Lemmas 3.4 and 3.5(a). By Theorem 3.3(a), it follows that S_{M_n} is the torus. In the case where one of B and W is empty we have (i) and (ii) implying

$$\dim[\text{Im}(c_{P_n} \circ c_{P_n})] = 0.$$

Therefore, in this case S_{M_n} is the plane, by Theorem 3.3(a).

Suppose that n is even. Now $i_{P_n}^2(x)$ is empty for every x in $\text{symb}(\overline{P}_n)$. Hence, we have the implications:

- (i') if $x \in B$, then $(c_{P_n} \circ c_{P_n})(x) = B$;
- (ii') if $x \in W$, then $(c_{P_n} \circ c_{P_n})(x) = W$.

It follows that if one of B and W is empty, then $\dim[\text{Im}(c_{P_n} \circ c_{P_n})] = 1$ and S_{M_n} is the projective plane. If both of B and W are non-empty, then

$\dim[Im(c_{P_n^\sim} \circ c_{P_n})] = 2$, and S_{M_n} is the Klein bottle. ■

Chapter 4

Transversals of Orientation – Reversing Circuits

4.A Preliminary Concepts and Relation with Imbalances

Given a finite collection Ω of subsets of a finite set X , a *transversal* of Ω is a subset T of X such that $T \cap Y$ is non-empty for every element Y of Ω . A *minimal* transversal is a transversal not properly contained in another transversal. A *minimum* transversal is a transversal of minimal cardinality. As usual, the adjectives minimal and minimum, as well as maximal and maximum, have this distinction, when applied to some concept.

This chapter establishes a minimax relation (Theorem 4.5), which includes in its proof an algorithm, polynomially bounded in the “size” of the map, to exhibit a minimum transversal of r -circuits in arbitrary projective maps.

Our aim in this first section is to relate transversals of r -circuits and imbalances (Lemma 4.1). Also we show (Proposition 4.2) that the problem of deciding whether a subset of eG_M is a minimum transversal of r -circuits in an arbitrary map M is *NP*-complete, in the sense introduced by Karp [13].

With the next lemma we switch from transversals of r -circuits to imbalances, which is technically convenient. Once more, because of the natural 1-1 correspondence between eG_M and $SQ(M)$, we do not distinguish explicitly their members or subsets.

(4.1) Lemma: For every map M , the minimal transversals of r -circuits in M^t are precisely the minimal imbalances in M .

Proof: Let I be a minimal imbalance in M . Denote by f_I the set of f_M -edges in the squares in I and let $\{A, B\}$ be the balancing partition inducing I . By the definition of balancing partition and of induced imbalance, f_I is precisely the set of edges with both ends in A or both ends in B . Hence, $\delta A = eC_M \setminus f_I = \bar{f}_I$. Therefore, we conclude the following equivalence: I is a minimal imbalance in M iff $\bar{f}_I$ is a maximal coboundary in C_M .

Proposition 4.1.1, proved below, states that a subset of edges in eG , for arbitrary map G , is a maximal coboundary in G if and only if its complement is a minimal transversal of odd circuits in G . Thus, applying Proposition 4.1.1, we conclude that, the complement f_I of a maximal $\bar{f}_I$, as above, is a minimal transversal of odd circuits in C_M . To conclude the proof it is enough to observe that I is a minimal transversal of r -circuits in M^t if and only if f_I is a minimal transversal of odd circuits in C_M . This equivalence follows from the definition of r -circuit. Provided Proposition 4.1.1 is established, the proof of the lemma is complete. ■

(4.1.1) Proposition: For an arbitrary graph G , a subset of edges in eG is a maximal coboundary in G if and only if its complement is a minimal transversal of odd circuits in G .

Proof: It is evident that the complement in eG of any coboundary in G is a transversal of odd circuits in G . Thus, to prove the proposition, it is enough to show that a minimal transversal of odd circuits is the complement in eG of a coboundary.

Let T be a minimal transversal of odd circuits in G . Hence the graph $G \setminus T$, obtained by deleting from G all the edges in T , is bipartite. Let $\{A, B\}$ be a bipartition of $G \setminus T$. We claim that every edge in T must have both ends in A or both ends in B . If edge $e \in T$ has one end in A and the other in B , then $T \setminus \{e\}$ is yet a transversal of odd circuits and so, T would not be a minimal transversal. This contradiction proves the claim. Therefore, the set of edges in the complement of T , each linking a vertex in A to a vertex in B , is the coboundary of A in G . ■

(4.2) Proposition: To determine whether a given subset of eG_M is a minimum transversal of r -circuits in a general map M is an NP -complete problem.

Proof: We show that the problem of determining whether a given subset of eG is a maximum coboundary in an arbitrary graph G is reducible to the problem stated in the proposition. Since the maximum coboundary problem is NP -complete, see Garey and Johnson [6], this is enough to establish (4.2). By Proposition 4.1, it is enough to reduce the maximum coboundary problem to the minimum imbalance problem.

Assume we are given a graph G for which we want to discover a maximum coboundary. Let us embed G arbitrarily in an orientable surface such as to obtain a t -map M^t with $G_M \cong G$. Observe that M is obtainable with an amount of work which is bounded by a polynomial in the number of edges of G . Specifically choose, for each vertex x of G , an arbitrary cyclic sequence of edges incident to x to go around x in M^t . These cyclic sequences form a v -ordering for M . Take the empty set to be the imbalance corresponding to this v -ordering. By Proposition 3.0 a unique map M is specified. Since the set of imbalances in a map M is a coset of the coboundary space of G_M , and since ϕ is an imbalance in M , we conclude that the set of imbalances in M is precisely the coboundary space of $G_M \cong G$. By Proposition 3.2.1, $\subseteq SQ(M)$ is an imbalance in M if and only if $SQ(M) \setminus T$ is an imbalance in $M^\sim$. Therefore, to find a maximum coboundary in G is the same as to

find a maximum imbalance in M , which is the same as to find a minimum imbalance in $M^\sim$. This concludes the proposition. ■

4.B Imbalances and Cycles in the Dual

In this section we prove that the imbalances of a map are cycles of a specific homology in the dual. We also prove that the set of minimal imbalances in a projective t -map is precisely the set of r -circuits in the dual t -map.

(4.3) Theorem: The set of imbalances in a map M is a class of homologous (mod 2) cycles in the dual D^t of M^t .

Proof: We already know that the set of imbalances in M is a coset of the coboundary space V of G_M . It follows that the symmetric difference of two arbitrary imbalances is a coboundary in G_M , and hence a boundary in D^t .

Therefore, to conclude the proof it is sufficient to establish that the set of squares in an imbalance I corresponds to a set of edges in G_D which is a cycle in G_D .

Let I be an arbitrary imbalance in a map M induced by a balancing partition $\{A_I, B_I\}$. Let us impose an orientation on the v_M - and the a_M -edges of C_M such that every v -gon of M becomes a directed polygon and such that each v_M -dart has tail in A_I and head in B_I .

Consider now the traversal of an arbitrary f -gon K , and an f_M -edge x in K . The two a_M -edges, preceding and following x , are traversed one in the same direction and the other in the opposite direction (according to the imposed orientation of the a_M -edges) if and only if the square containing f_M -edge x is in the imbalance I . Observe that in the complete traversal of K the number of times that there is a reversal of direction when we go from an a_M -edge to the next is even. Therefore, the number of squares in I met by the traversal of K is even. Thus, the number of squares in I met once by

the traversal of K is even. Observe that this set of squares corresponds to a coboundary of the vertex K in G_D (the vertices of G_D are f -gons). Since K is arbitrary, the imbalance I is orthogonal to every element in the coboundary space F of G_D . That is to say, I belongs to $F^\perp$ the cycle space of G_D . This establishes the theorem. ■

Let M be a projective map. By Lemma 2.6 and the previous theorem, the set of imbalances in M is either the set V of boundaries in D^t or a set of the form $x + V$, where x is an r -cycle in D^t . The set V of coboundaries in G_M is discarded, since V is the set of imbalances in M iff M is orientable, which is not the case by Theorem 1.7. Therefore, we obtain the following lemma:

(4.4) Lemma: The minimal imbalances in a projective map M are precisely the r -circuits in G_D where D is the dual of M .

Proof: We proved that the set of imbalances in M is of the form $X + V$, where x is an r -cycle in G_D and V is the coboundary space of G_M . The proof of the lemma follows, then, by the definition of r -circuit as a minimal r -cycle. ■

4.C Main Theorem

A collection Q of subsets of X is a *disjoint* collection if the members of Q have no elements in common. The main theorem of the chapter is the following:

(4.5) Theorem: For an arbitrary projective map M with G_M eulerian, a minimum transversal of r -circuits in M^t is equal in cardinality to a maximum disjoint collection of r -circuits in M^t .

By Lemma 4.1, the minimal transversals of r -circuits of a map M are precisely the minimal imbalances of M . Hence, in the statement of (4.5) we

can replace “transversal of r -circuits in M^t ” by “imbalances in M ”.

As we show in the sequel, the hypothesis that G_M is eulerian is not essential, and the algorithmic proof of (4.5) provides us with a polynomial algorithm to determine the size of a minimum imbalance in general projective maps. As we showed in Proposition 4.2, the minimum imbalance problem is NP -complete. It is, therefore, of interest to know exactly for which classes of maps (4.5) holds. In our conclusions of Chapter 5 we present another class of maps in which (4.5) holds.

By Lemma 4.4, the minimal imbalances in a projective map M are precisely the r -circuits of G_D , where D is the dual of M . This provides us with yet another equivalent form of statement 4.5: we can replace “transversal of r -circuits in M^t ” by “ r -circuit in D^t ”, where D is the dual of M .

It is this form of (4.5), making use of r -circuits in the dual, which is suitable for the proof that we present in Section 4. Before we introduce, in the next section, the concept of “reducer”, which is the most important concept in the proof of (4.5) we give a corollary avoiding the hypothesis that G_M is eulerian.

Given a map M let us denote by M_d the map which corresponds to the t -map M^t in which each edge of G_M is replaced by a pair of edges forming a bounding digon in the surface S_M . Let D_d be the dual of M_d . It is evident that each circuit in G_{D_d} corresponds to a circuit in G_D half as long in length. In particular, each r -circuit in D^t corresponds to an r -circuit in D_d^t half as long in length. Therefore, the cardinality of a minimum r -circuit in D_d^t is twice the cardinality of a r -circuit in D^t . Consequently, the cardinality of a minimum transversal of r -circuits in M_d^t is twice the cardinality of a minimum transversal of r -circuits in M^t . This observation implies the following

(4.5) Corollary: For an arbitrary projective map M , a minimum transversal of r -circuits in M^t has cardinality equal to half the cardinality of a maximum collection of r -circuits in M^t with the property that each edge of G_M is used at most twice in the collection.

Proof: The corollary follows from the application of Theorem 4.5 to the map M_d , describe above. ■

In the next section we introduce the concept of “reducers” The elimination of reducers is the basis for a polynomial algorithm to establish the minimax results stated. In Figure 4.1 below, we show the result of the application of this algorithm, illustrating the Corollary of 4.5. Figure 4.1 presents a projective t -map M^t , for which a minimum transversal of r -circuits has cardinality 5: the set of five edges which appear into two pieces is clearly a transversal of r -circuits in M^t . The proof that the set of these five edges is a minimum transversal is established by exhibiting ten r -circuits in which each edge is used at most twice by the ten circuits.




Figure 4.1

4.D Splitting Reducers

A pre-requisite for the definition of a reducer is the notion of contractibility. The definition that we give is suitable for our purposes. This becomes apparent in Corollary 4.8.1.

If R is a subset of faces of M^t , we denote by eR the set of edges of G_M that appear (one or two times) in the facial paths of the faces in R .

A subset R of faces of M^t is contractible if its boundary is a circuit and if every circuit in G_M included in eR is a boundary in M^t .

Observe that R is contractible iff the union of the faces in R together with the (topological) subgraph of G_M generated by eR is homeomorphic to a closed disc.

We call the attention of the reader to the fact that for a subset of faces R whose boundary is a circuit it is not the case, in general, that either R or its complement $\overline{R}$ is contractible. However, it is the case if M is projective, as we establish later in Corollary 4.8.1. In the Klein bottle we may have a subset R of faces whose boundary is a circuit but such that neither R nor $\overline{R}$ is contractible.

For a map M in which G_M is eulerian, we denote by $SP(M^t)$ the set of smooth paths of M^t .

For a map M in which G_M is eulerian, a non-empty subset R of faces of M^t is called a *reducer* for M^t if R is contractible and the circuit which constitutes the boundary of R is induced by at most two subpaths of members of $SP(M^t)$.

A *type 0 reducer* is a reducer whose boundary is induced by a (complete) smooth path. Note that a type 0 reducer cannot be minimal: if R_0 is a type 0 reducer there is a reducer R , (not necessarily of type 0) such that R is a subset of faces properly contained in R_0 .

A *type 1 reducer* is a reducer whose boundary is induced by a unique proper reentrant subpath of a smooth path.

A *type 2 reducer* is a reducer whose boundary is induced, not only by one but by two subpaths of smooth paths with the same ends.

Observe that a reducer R of type 1 which is also a minimal reducer, in the sense that R does not properly contain a subset of faces S which is also a reducer of type 0, 1 or 2, is necessarily a singleton $\{f\}$ where f is a face whose boundary is a loop in G_M (monovalent face).

We identify now an important property of any minimal reducer R of type 2. Denote by p and q the two subpaths of members of $SP(M^t)$ which induce the boundary of R . A *segment* of R is a maximal subpath of a member of $SP(M^t)$ with at least one edge whose edges are contained in $eR \setminus \partial R$, where ∂R , recall, is the boundary of R . It is evident that R has the following property TRANS: for an arbitrary segment s , of R , one end of s is an interior vertex of p and the other is an interior vertex of q .

A *transreducer* is either a monovalent face or else a type 2 reducer satisfying property TRANS, described above. The transreducers are important for two reasons: first, given a reducer R it is algorithmically simple to find a transreducer S contained in R ; second, property TRANS in a reducer is enough for our purposes, which are described in the sequel.

Figure 4.2 below shows examples of transreducers.




Figure 4.2

An in a reducer R it a vertex which is a common end of the path or paths which induce the boundary of R . Hence, a reducer has zero, one or two angular points, according as to whether it is of type 0, type 1 or type 2.

The splitting of a reducer R (not of type 0) is a local topological operation effected in M^t near an angular point v of R , defined as follows. Denote by e_1 and e_2 the two edges in $\partial(R)$ which are incident to v . Let f_1 and f_2 be the opposite edges to e_1 and e_2 at v , respectively. Separate the topological paths formed by f_1 and e_2 from the one formed by f_2 and e_1 , so that both paths miss v . If v has valency 4, then it is replaced by two bivalent vertices after the splitting. See Figure 4.3.




Figure 4.3

Let us denote by N^t the t -map obtained from M^t by the splitting of a transreducer. Let E be the dual of N . It is evident that the classes of homology in $CS(G_D)/V_M$ and in $CS(G_E)/V_N$ are in natural 1-1 correspondence. We use this correspondence to identify those classes and to state the following important

(4.6) Lemma: For a map M with G_M eulerian, the splitting of a transreducer does not change the cardinality of a minimum circuit in a non-null homology class of D^t , where D is the dual of M .

Proof: To prove this lemma, consider a non-bounding circuit c in E^t , where E^t is the dual of N^t , the t -map obtained from M^t by the splitting of

a transducer R . If c is also a circuit in G_D , there is nothing to be proved. Therefore, we may consider that the polygon corresponding to c has among its vertices some which are new (triangular) faces in N^t introduced by the splitting. See Figure 4.4. (In the case that the angular point of R is 4-valent, then two faces of M^t becomes a (new) face of N^t .)




Figure 4.4

Let f_1 be the face of M^t incident to the two edges not in the boundary of R which are incident to the angular point of R near which the splitting is effected. If c is not a circuit in the dual graph, G_D' then the polygon corresponding to c has an a_1 (a graph induced by a path without repeated vertex-terms) which goes from vertex f_1 uses some new vertices (new faces in N^t) and finishes in f_2 which is the first old face not in R .

We claim that arc a_1 can be replaced by an arc a_2 which also links f_1 to f_2 has the same number of edges as a_1 , and does not have among its vertices, either new faces of N^t or faces in R . See Figure 4.4. The new arc a_2 in G_E starts at f_1 , crosses an edge of G_N which corresponds to an edge in the extension of the boundary of R incident to the angular point, crosses some edges of G_N which have a vertex incident to the boundary of R , until it reaches f_2 . The crossing points between the (topological) paths

corresponding to a_1 and a_2 and the topological edges in G_E are evidently in 1-1 correspondence with eG_E and eG_N . Denote by d_1 the set of edges of a_i , $i = 1, 2$. Observe that $d_1 \subseteq c$, $d_2 \cap c = \emptyset$, and that $d_1 + d_2$ is a boundary in E^t . Observe also that $c' = (c \setminus d_1) + d_2$ is a circuit in G_N as well as in G_E , and it is homologous to c . By the fact that c is minimum in its homology class, $|c'| \geq |c|$, that is $|d_2| \geq |d_1|$. It is evident by construction, that a_2 links f_1 to f_2 and does not have among its vertices either new faces or faces in R . To conclude the claim, we show that the number of crossings of a_2 with edges in G_N , i.e., $|d_2|$, is at most equal to the number of crossings of a_1 with edges in G_N , i.e., $|d_1|$. The first crossing of a_2 is compensated for by the last crossing of a_1 , with an edge in the boundary of R . A crossing between a_2 and an edge in the extension of a segment of a smooth path whose edges are contained in eR , is compensated for, since R satisfies TRANS, by an odd number of crossings between a_1 and edges in the same segment. Thus $|d_2| \leq |d_1|$, proving that $|c'| = |c|$. Thus, we have shown that the minimum circuit in a non-null homology class of D^t is at most equal in cardinality to a minimum circuit in the corresponding homology class of E^t . The converse inequality is trivial, because any circuit in G_D is also a circuit in G_E , (with the obvious identifications of edges we have $eG_D \subseteq eG_E$), and the homologies classes are maintained. This concludes the proof of the lemma. ■

It is possible that the minimum cycle of a given homology class is not a circuit. For this reason we include the following strengthening of Lemma 4.6.

(4.7) Theorem: For a map M with G_M eulerian, the splitting of a transducer does not change the cardinality of a minimum cycle in a non-null homology class of D^t , where D is the dual of M .

Proof: Let N and E be the pair of dual maps corresponding to M and D after the splitting is performed. Let c be a minimum cycle in a non-null

member H of $CS(G_E)/V_N$. Let us decompose c into pairwise disjoint circuits

$$c = c_1 + c_2 + \cdots + c_k.$$

Each circuit c_i is not a member of V_N , the coboundary space of G_N ; otherwise $c \setminus c_i$ would be a cycle in H with fewer edges than c , contradicting the assumption.

By Lemma 2.6, for each circuit c_i in G_E which is not a circuit in G_D , we can obtain a circuit c'_i , homologous to c_i , of the same cardinality as c_i , and which is a circuit in G_D . Observe that, since $c' = (c \setminus c_i) + c'_i$ is a cycle in G_E , the minimality of $|c|$ in H implies that $c_j \cap c'_i = \emptyset$, for $j \neq i$. Hence, $|c'| = |c|$. By replacing in this way each c_i which is not in G_D by a corresponding c'_i we get a cycle d , homologous to c , of the same cardinality as c , and which is a cycle in G_D . Thus, we have shown that the minimum cycle in the homology class of D^t identified with H is equal in cardinality to a minimum cycle in H . The converse inequality is trivial, because any cycle in G_D is also a cycle in G_E , and the homologies are maintained. This concludes the proof of the theorem. $\blacksquare$

With the above theorem we end this section. In the next section we particularize it to projective maps so as to prove the main result of the chapter, Theorem 4.5.

4.E Proof of the Main Theorem

In this section we give a proof of Theorem 4.5. Before, however, we establish some lemmas. The following lemma is implicit in the proof of Theorem 2.14. Now we include it as an explicit statement.

(4.8) Lemma: For every map M , the maximum number of pairwise disjoint odd polygons in C_M is at most $\xi(M)$.

Proof: By Proposition 2.6, applied to t -map (C_M, S_M) , the quotient space

$$\frac{CS(C_M)}{\langle SQ(M) \rangle + \langle vG(M) \rangle + \langle fG(M) \rangle}$$

has exactly $\xi(M)$ elements (classes of homology). By Theorem 1.8, no subset of circuits induced by pairwise disjoint odd polygons is in the space generated by the set of edges in the squares, v -gons, and f -gons of M . Hence, the set of edges in each polygon of an arbitrary disjoint collection of odd polygons in C_M represents a distinct class of homology. ■

(4.8.1) Corollary: Assume that M is a projective map and R is a subset of faces in M^t whose boundary is a circuit. Then, either R or $\overline{R}$, the complement of R in the set of faces of M^t , is contractible.

Proof: Assume indirectly that both R and $\overline{R}$ are not contractible. By definition, this implies that there exist non-bounding circuits S_1 and S_2 (circuits which are not boundaries), for which $S_1 \subseteq eR$, $S_2 \subseteq e\overline{R}$. By Proposition 3.2.2., S_1 and S_2 are r -circuits. Since S_1 and S_2 can meet only along the common boundary of R and $\overline{R}$, it follows that we can find disjoint odd (in cardinality) circuits, S'_1 and S'_2 , in C_M such that $\psi_c^M(S'_i) = S_i$, $i = 1, 2$. But this contradicts lemma 4.8, since it implies the existence of two disjoint odd polygons in C_M , and $\xi(M) = 1$. ■

For disjoint cycles S_1 and S_2 in G_M we say that S_1 and S_2 *cross* if for every pair (S'_1, S'_2) of cycles in C_M such that $\psi_c^M(S'_1) = S_1$ and $\psi_c^M(S'_2) = S_2$, $S'_1 \cap S'_2 \neq \emptyset$.

This formal definition of crossing is a set-theoretical description of a crossing at a vertex of two edge-disjoint polygons in M^t .

(4.8.2) Corollary: For any projective map M , if S_1 and S_2 are disjoint r -cycles in M^t , then S_1 and S_2 cross.

Proof: Assume that S_1 and S_2 do not cross. Thus, there exists disjoint odd cycles S'_1 and S'_2 in C_M such that $\psi_c^M(S'_i) = S_i$, $i = 1, 2$. Therefore, there exists disjoint odd polygons Q_1 and Q_2 in C_M such that $eQ_i \subseteq S'_i$. But this contradicts lemma 4.8, because $\xi(M) = 1$. ■

A path π is *simple* if no vertex appears twice as vertex-term in π . A path π is *semi-simple* if it is simple or if it is reentrant and the only vertex which appears twice as vertex-term in π is its origin which is the same vertex as its terminus.

The technique used to prove 4.5 is to effect the splitting of transreducers. To find a transreducer it is enough to be able to locate reducers. To locate a reducer we first find a pre-reducer, defined as follows.

A pre-reducer in a t -map M^t with G_M eulerian is an ordered pair (π_1, π_2) of subpaths of members of $SP(M^t)$, such that π_1 is semi-simple, π_2 is non-degenerate, π_1 and π_2 have only their ends as common vertices, and $e\pi_1 \cup e\pi_2 = e\pi_1 + e\pi_2$ is an e -cycle in M^t .

We remark that in a pre-reducer (π_1, π_2) , π_1 , may be degenerate, being formed by just one vertex.

A *straight* pre-reducer (π_1, π_2) is a pre-reducer in which π_2 is semi-simple.

(4.9) Proposition: For a projective map M with G_M eulerian, if (π_1, π_2) is a straight pre-reducer, then there is a reducer R for M^t such that $\partial(R) = e\pi_1$ or $\partial(R) = e\pi_1 + e\pi_2$.

Proof: Assume that π_1 and π_2 are both reentrant. By definition of straight pre-reducer, $e\pi_1$ and $e\pi_2$ are both r -circuits or both s -circuits. Observe, however, that $e\pi_1$ and $e\pi_2$ do not cross. Hence, by Corollary 4.8.2, they are both s -circuits. By Proposition 3.2.2, $e\pi_1$ is a boundary of a subset X of faces. By Corollary 4.8.1, either X or its complement $\overline{X}$ is contractible. Let R be a contractible member of $\{X, \overline{X}\}$. By definition, it follows that R is a reducer, and since $\partial R = e\pi_1$, the proposition follows.

Assume now that π_1 (and π_2) are not reentrant. By definition of straight pre-reducer $e\pi_1 + e\pi_2$ is an s -circuit. Applying Proposition 3.2.2 and Corollary 4.8.1 as before, with $e\pi_1 + e\pi_2$ in the place of $e\pi_1$, we define a reducer R such that $\partial R = e\pi_1 + e\pi_2$ concluding the proof. ■

(4.10) Lemma: For a projective map M with G_M eulerian, if (π_1, π_2) is a pre-reducer which is not straight, then there exists a pre-reducer (π_3, π_4) such that $e\pi_3 \cup e\pi_4$ is a proper subset of $e\pi_1 \cup e\pi_2$.

Proof: Since π_2 is not semi-simple, we can write $\pi_2 = \alpha \circ \beta \circ \gamma$, where $e\beta$ is a circuit. If $e\beta$ is an s -circuit in M^t , then take π_4 to be the degenerate path whose only vertex is the origin of the reentrant path β , and take π_3 to be β ; it follows that (π_3, π_4) satisfies the condition of the lemma. Therefore, we may assume that $e\beta$ is an r -circuit in M^t . Since $e\pi_1 + e\pi_2$ is an s -cycle and since $e\pi_2 = e\alpha + e\beta + e\gamma$, it follows that $e\pi_1 + e\alpha + e\gamma$ is an r -cycle. By Corollary 4.8.2, $e\pi_1 + e\alpha + e\gamma$ crosses $e\beta$.

It follows that at least one of the following vertex-terms must exist:

- (i) a first vertex-term of α^{-1} other than its origin which is also a vertex-term of β ;
- (ii) a first vertex-term of γ other than its origin which is also a vertex-term of β .

The situation is symmetric, up to taking inverse paths, and so we may consider that (ii) holds. Denote by v the vertex-term described in (ii), and by γ_1 the proper subpath of γ that goes from its origin up to v . Let us write β as $\beta_1 \circ \beta_2$ where the terminus of β_1 and the origin of β_2 coincide with v . One of β_1 and β_2 may be degenerate. The reentrant paths γ_1, β_1 , and β_2^{-1} all have the same ends. Since $e\beta_1 + e\beta_2$ is an r -cycle (indeed an r -circuit), it follows that $e\beta_1 + e\gamma_1$ is an r -cycle iff $e\beta_2 + e\gamma_1$ is an s -cycle. By the symmetry of the situation, we may assume that $e\beta_1 + e\gamma_1$ is an e -cycle. We can now easily verify that (β_1, γ_1) is a pre-reducer. Since $e\beta$ is a circuit, β_1

is semi-simple. By definition of v , γ_1 is non-degenerate and β_1 and γ_1 have in common only their ends. Since $e\beta_1 + e\gamma_1$ is an s -cycle, it follows that (β_1, γ_1) is a pre-reducer. Put $\pi_3 = \beta_1, \pi_4 = \gamma_1$, and the proof of the lemma is complete. ■

The following lemma is the last one before we proceed to the proof of Theorem 4.5.

(4.11) Lemma: For a projective map M with G_M eulerian, if $\pi \in SP(M^t)$ is such that $e\pi$ is an s -cycle in M^t , or if $\pi \in SP(M^t)$ is not semi-simple, or if two elements of $SP(M^t)$ have more than one vertex-term in common, then there exists a pre-reducer in M^t .

Proof: Suppose that $\pi \in SP(M^t)$ is such that $e\pi$ is an s -cycle in M^t . Take π_0 to be a degenerate path whose only vertex term is a vertex-term of π . By definition, (π_0, π) is a pre-reducer in M^t .

Assume next that every member of $SP(M^t)$ induces an r -cycle and that $\pi \in SP(M^t)$ is not semi-simple. Let us write π as $\alpha \circ \beta$, where $e\beta$ is a circuit. (In particular this implies $|e\beta| \geq 1$.) If $e\beta$ is an s -circuit, then consider δ to be the degenerate path whose only vertex is the origin of β . The pair (δ, β) is a pre-reducer in M^t . Thus, we may assume that $e\beta$ is an r -circuit. But since $e\alpha + e\beta = e\pi$ is an r -cycle, it follows that $e\alpha$ is an s -cycle. Take δ to be the degenerate path whose only vertex-term is the origin of α . The pair (δ, α) is a pre-reducer in M^t .

Finally, assume that for every $\pi \in SP(M^t)$ $e\pi$ is an r -cycle and π is semi-simple. Suppose that π_1 and π_2 have two distinct vertices v and w as common vertex-terms. Write π_1 as $\alpha \circ \beta^{-1}$, π_2 as $\gamma \circ \delta^{-1}$, where α, β, γ and δ all have origin v and terminus w . If $\alpha + \gamma^{-1}$ induces an r -cycle, then $\alpha \circ \delta^{-1}$ must induce an s -cycle, because $e\gamma + e\delta = e\pi_2$ is an r -cycle, and $e\alpha + e\gamma = e\alpha + e\delta + e\pi_2$. Therefore, we may suppose, without loss of generality, that $e\alpha + e\gamma$ is an s -cycle. (If not, take $e\alpha + e\delta$ in place of $e\alpha + e\gamma$.) Let u be the first vertex-term of path α such that u is also a vertex-term of

γ . Vertex-term u must exist because w is common to both α and γ . Let α_1 be the subpath of α which goes from v until u . Let γ_1 be the subpath of γ which goes from v until u . We want to show that (α_1, γ_1) is a pre-reducer in M^t . By definition of u , α_1 is semi-simple, $|e\gamma_1| \geq 1$, and paths α_1 and γ_1 have only their ends as common vertices. To conclude that (α_1, γ_1) is a pre-reducer, it is enough to show that $e\alpha_1 + e\gamma_1$ is an s -cycle. Suppose that $e\alpha_1 + e\gamma_1$ is an r -cycle. Let α_2 and γ_2 be defined by the equations $\alpha = \alpha_1 \circ \alpha_2$, $\gamma = \gamma_1 \circ \gamma_2$. Observe that the cycle $e\alpha_2 + e\gamma_2$ is an r -cycle, since $e\alpha_1 + e\gamma_1$ is an r -cycle and $e\alpha + e\gamma$ is an s -cycle. Observe also that, again by the definition of u , $e\alpha_1 + e\gamma_1$ does not cross $e\alpha_2 + e\gamma_2$. But this contradicts Corollary 4.8.2. Therefore, $e\alpha_1 + e\gamma_1$ is an s -cycle and so, (α_1, γ_1) is a pre-reducer, concluding the proof of the lemma. ■

Now we are in a position to prove Theorem 4.5 in the following equivalent form, involving r -circuits in the dual:

(4.5 restated) Theorem For an arbitrary projective map M with G_M eulerian, a minimum r -circuit in D^t is equal in cardinality to a maximum disjoint collection of r -circuits in M^t .

Proof: The proof of the theorem is divided into two parts. In the first part we show how to reduce the proof of the general case to the special case in which M^t is a system of projective lines, which we define next. In the second part we give a proof for this special case.

A projective t -map M^t with G_M eulerian is called a *system of projective lines* if every member of $SP(M^t)$ induces an r -circuit and for two arbitrary members, π_1 and π_2 of $SP(M^t)$ the polygons defined by π_1 and π_2 have precisely one vertex in common.

Now we show how to reduce the Proof of 4.5 from an arbitrary M^t with G_M eulerian to a system of projective lines. Suppose we are given a projective t -map M^t with G_M eulerian. If there exists a transreducer R for M^t , then Lemma 4.6 shows that the splitting of R does not change the cardinality of

a minimum r -circuit in the dual. Moreover, a disjoint collection of r -circuits in the t -map after splitting corresponds to a disjoint collection of r -circuits in the original t -map before splitting. Therefore, a proof of the theorem for the split t -map implies a proof for the original t -map.

We now introduce a measure relative to which the t -map obtained by splitting a transreducer is a smaller t -map. For a t -map M^t , denote by $\mu(M^t)$ the ordered pair of non-negative integers $(\lambda(M^t), f(M^t))$, where,

$$\lambda(M^t) = \sum_{v \in VG_M} \max\{0, \text{val}_{G_M}\{v\} - 4\},$$

and $f(M^t)$ is the number of faces of M^t . We say that $\mu(N^t) < \mu(M^t)$ if $\lambda(N^t) < \lambda(M^t)$ or if $\lambda(N^t) = \lambda(M^t)$ and $f(N^t) < f(M^t)$. (Lexicographic order.)

Observe that if N^t is obtained from M^t by splitting a reducer, then $\mu(N^t) < \mu(M^t)$ this follows because if v , the angular point near which the splitting is performed, has valency greater than 4, then $\lambda(N^t) < \lambda(M^t)$; if v has valency 4, then $\lambda(N^t) = \lambda(M^t)$, and $f(N^t) = f(M^t) - 1$. This implies that the number of successive splittings of transreducers that we can perform in a t -map M^t before reaching a t -map, N^t free of transreducers is at most $2|eG_M| + f(M^t)$.

The above analysis implies that we may restrict ourselves to prove 4.5 for maps free of transreducers. We now show that a map free of transreducers is system of projective lines. If M^t is free of transreducers, M^t is free of reducers, because a reducer always contains a transreducer. Since a straight pre-reducer implies a reducer, by Proposition 4.9, it follows that M^t has no straight pre-reducer. By Lemma 4.10 a pre-reducer which is not straight implies a “smaller” pre-reducer. Hence, any pre-reducer implies a straight pre-reducer. As a consequence, M^t is free of pre-reducers. By lemma 4.11, this implies that:

- (i) for $\pi \in SP(M^t)$, $e\pi$ is an r -cycle;
- (ii) if $\pi \in SP(M^t)$, then π is semi-simple;

- (iii) if π_1 and π_2 are distinct members of $SP(M^t)$, then π_1 and π_2 have at most one vertex-term in common.

Items (i) and (ii) imply that if $\pi \in SP(M^t)$, then $e\pi$ is an r -circuit. Item (iii) implies that the two polygons induced by two arbitrary members of $SP(M^t)$ have at most one vertex in common. By Corollary 4.8.2, two r -circuits must cross. Therefore, every pair of polygons as above has precisely one vertex in common. This proves that if M^t is free of transreducers, then M^t must be a system of projective lines. The first part of the proof is complete.

Now we conclude the theorem for a system of projective lines M^t . Explicitly, we exhibit a disjoint collection Ω of r -circuits in M^t and an r -circuit R in D^t such that $|\Omega| = |R|$. The collection Ω is simply $\{e\pi : \pi \in SP(M^t)\}$. Let $n = |\Omega|$. To obtain R , let p be the closed topological path corresponding to an arbitrary member of $SP(M^t)$ and take a point P near p in S_M . Refer to Figure 4.5 for this description. Proceed from P parallel to p until the point Q opposite to P relative to p is reached. Close the topological path p' which we are describing by linking Q to P , crossing p . The closed topological path p' which we described corresponds to an r -circuit R in D^t . The crossings of p' with the points in G_M are in 1-1 correspondence with the edges in R . Each crossing of p' , with the exception of the last described, corresponds to a crossing of p with another member of $SP(M^t)$; hence, they are $n - 1$ in number. Therefore, $|R| = n = |\Omega|$, proving the Theorem. ■




Figure 4.5

To conclude this section we show with an example that even for the torus the maps free of reducers can be quite complicated. Consider, for example, the toroidal t -map free of reducers shown in Figure 4.6. There are three smooth paths as pointed by different thickness on the edges of the illustration. Any two such smooth paths meet each other seven times, illustrating the point that straight pre-reducers do not imply reducers in general.




Figure 4.6

4.F Algorithmic Considerations

In this section we present an algorithm, PROJMINMAX, which has as its input a projective map M_0 with G_{M_0} eulerian and as its output a pair (Ω_0, R_0) , defined as follows:

- (i) Ω_0 is a collection of cyclic paths in G_{M_0} , such that the corresponding cycles in M^t form a disjoint collection of r -circuits.
- (ii) R_0 is an r -circuit in G_{D_0} , where D_0 is the dual of M_0 .

Moreover, the pair (Ω_0, R_0) satisfies $|\Omega_0| = |R_0|$. This is a pair whose existence is guaranteed in theorem 4.5.

We also show that the amount of space (space complexity) and the number of basic steps (time complexity) of PROJMINMAX are bounded by polynomials n , the “size” of the map, where $n = |eG_{M_0}| = |SQ(M_0)|$.

We say that an algorithm A has *time (space) complexity* $O(n^k)$, if there exists a constant C such that the number of basic steps (bytes of memory required) in the execution of A to solve a problem whose input has size n is at most Cn^k , except for finitely many values of n .

The algorithm that we present is suggested by the theory for finding transreducers and by the inductive step consisting in splitting a transreducer, which form the basis for the proof of Theorem 4.5. However, in the algorithmic realization, we do not actually split transreducers.

In order to avoid splittings in PROJMINMAX we introduce the following Ω -parametrized concepts, where Ω is an edge-disjoint collection of cyclic paths in G_M whose edge-sets partition eG_M , for map M with G_M eulerian. The following concepts generalize the ones already described in which Ω is defined by the smooth paths of M^t .

An Ω -*reducer* for M^t is a subset of faces R such that R is induced by at most two subpaths of members of Ω and such that eR contains only bounding circuits.

If R is an Ω -reducer for M^t , then a *segment* of R is a maximal subpath of a member of Ω with at least one edge, whose edges are contained in $eR\partial R$.

An *angular point* in an Ω -reducer R is a vertex in which the extension of the path or paths forming the boundary of R cross.

Property *TRANS* in a Ω -reducer R with two angular points and boundary induced by two subpaths p and q of members of Ω is defined as follows: for every segment s of R , the extension of s in Ω crosses p and not q at one end of s and q and not p at the other end.

An Ω -*transreducer* is an Ω -reducer R with two angular points for which property TRANS holds or else an Ω -reducer R with only one angular point that is formed by a single face.

An Ω -pre-reducer is an ordered pair (π_1, π_2) of sub-paths of members of Ω such that π_1 does not self-cross, π_2 is non-degenerate, π_1 and π_2 cross only at their ends (when extended), and $e\pi_1 \cup e\pi_2 = e\pi_1 + e\pi_2$ is an s -cycle in M^t . An Ω -pre-reducer is *straight* if π_2 does not self-cross.

The proof of the three following “ Ω -extensions” are repetitions of the ones given in 4.9, 4.10, and 4.11, where $SP(M^t)$ takes the role of Ω .

(4.9)* (Extension of 4.9) For projective map M with G_M eulerian and collection Ω of cyclic paths whose edge-sets partitions eG_M , if (π_1, π_2) is a straight Ω -pre-reducer, then there is an Ω -reducer R for M^t such that $\partial R = e\pi_1$ or $\partial R = e\pi_1 + e\pi_2$. ■

(4.10)* (Extension of 4.10) For (M, Ω) as above, if (π_1, π_2) is an Ω -pre-reducer which is not straight, then there exists an Ω -pre-reducer (π_3, π_4) such that $e\pi_3 \cup e\pi_4$ is a proper subset of $e\pi_1 \cup e\pi_2$. ■

(4.11)* (Extension of 4.11) For a pair (M, Ω) as above, if $\pi \in \Omega$ is such that $e\pi$ is an s -cycle in M^t , or if $\pi \in \Omega$ self-crosses, or if two elements of Ω cross more than once, then there exists an Ω -pre-reducer in M^t . ■

In Figure 4.7 we give the flow-graph for PROJMINMAX. It operates on the map $M = (C_M, V_M, f_M)$ rather than on M^t . The collection Ω at every stage of the algorithm is a collection of cyclic paths free of repeated edges whose edge-sets partition eG_{M_0} . We observe that a convenient way to keep track of Ω is, for every v -gon at M_0 , to specify a pairing in its v_M -edges. This pairing is possible, since G_{M_0} is eulerian. Given Ω , we define this pairing as follows. For each subsequence of a member of Ω of the form (e_1, V, e_2) , where e_1 and e_2 are edges of G_M and v is a vertex of G_M , denote by E_1 and E_2 the M -squares that correspond to edges e_1 and e_2 ; denote by v the v -gon of M which corresponds to the vertex v . We say that the v_M -edge which is in E_1 and in V and the v_M -edge which is in E_2 and in V are Ω -mates. Observe that the complete collection of pairs consisting of Ω -mates is enough to specify Ω .

itself.

Splitting a transreducer is replaced by the suitable change of two pairs of Ω -mates: $\{a, b\}, \{c, d\}$ become $\{a, c\}, \{b, d\}$ or $\{a, d\}, \{c, b\}$. Observe that, in this way, Ω changes but not the map M , as would be the case if splittings were performed.




Figure 4.7: Flow-graph for PROJMINMAX

In the proof of 4.5 restated, we show that the number of splittings is bounded by $2|eG_M| + f(M_0) \leq 0(n)$, where $f(M_0)$ is the number of f -gons in M_0 . Modifying Ω -mates is a suitable way to perform splittings; it follows that $0(n)$ is the order of the number of times that we answer the question “Is there an Ω -pre-reducer P ?”. See Figure 4.7.

For each positive answer of the above question, we answer in the negative at most $0(n)$ times the following question: is P straight? This follows from (4.10)*, and from the fact that an Ω -pre-reducer can get smaller at most n times.

For each positive answer to the first question in Figure 4.7, only once is the second question to be answered in the positive. For each such event, the third question “Is T an Ω -transreducer?” is answered at most $0(n)$ times. This is so because a test to decide whether TRANS is satisfied ends in a negative answer only if a smaller *Omega*-reducer is found. Hence, this can happen at most n times.

From the above analysis we conclude that the second and third questions are answered in the execution of PROJMINMAX at most $0(n^2)$ times and that the first question is answered at most $0(n)$ times.

We use (4.11)* to bound the time complexity for an answer to the first question by $0(n^2)$. The specific recipe on how to decide whether there exists an Ω -pre-reducer is given by an adaption of the proof of (4.11) for the corresponding Ω -extension (4.11)*. The bound $0(n^2)$ is sufficient for the decision.

An answer for the second and third questions can be obtained in $0(n)$ basic steps. To find a smaller Ω -pre-reducer, refer to (4.10)*, modelling its proof according to the proof of (4.10). This procedure is seen to take at most $0(n)$ steps. To find an Ω -reducer from a straight Ω -pre-reducer it also takes $0(n)$ steps. The same is true for finding a smaller Ω -reducer, in the case that the Ω -reducer at hand does not satisfy TRANS.

When an Ω -transreducer T is identified, we suitably change the two pairs of Ω -mated v_M -edges, an operation which corresponds to split T . This is followed by placing T (as a set of f -gons, for instance) in the top of a stack called TSTACK. This structure is important for the revision of R (explained below) in the second phase of the algorithm. The time complexity of the above two operations is each at most $0(n)$. The space required to keep all the structures so far is dominated by the space to keep TSTACK, which takes at most $0(n^2)$ bytes of memory.

This analysis implies that the time complexity for the first part of PROJMINMAX, which terminates with a negative answer to the first question, is $0(n^3)$.

The symbol R represents throughout a circuit in C_M . Given Ω , the Ω -weight of R is defined as the number of Ω -mated pairs with the property that exactly one member of the pair is a v_M -edge in R . The Initial R is any circuit in c_M with the property that $\psi_c^M(R)$ is the circuit induced by an arbitrary member of Ω_0 and such that its Ω_0 -weight is $|\Omega_0|$. See Figure 4.5. When we change Ω (crossing), the Ω -weight of R can increase by 2. The purpose of the revision of R is to find another R with Ω -weight equal to $|\Omega_0|$. Given the Ω -transreducer, this revision is a slight adaptation in terms of C_M of the Proof of Lemma 4.6. Refer to Figure 4.4. The v_M -edges of C_M correspond precisely to the crossings presented in this figure. The question asking whether TSTACK is empty is made at most $0(n)$ times. Instruction $T = \text{TOP}(\text{TSTACK})$ is self-explanatory; instruction $\text{POP}(\text{TSTACK})$ removes the topmost element of TSTACK. Each individual procedure of the second phase has complexity $0(n)$. Therefore, the overall execution of the second phase of PROJMINMAX has complexity $0(n^2)$. The space complexity of this phase is also $0(n^2)$ since TSTACK requires more memory than everything else.

When TSTACK is empty, we have that $\Omega = SP(M^t)$, and that the Ω -weight of the current R is equal to $|\Omega_0|$. For every pair of f_M -edges in R incident to a v -gon, we adjust R so as to use the shorter of the two possibilities to go around the v -gon. This does not change the Ω -weight of R . Observe

that, since Ω is now $SP(M_0^t)$, the Ω -weight of R , which is $|\Omega_0|$, is equal to the number of v_M -edges in R . But this is the cardinality of $R_0 = \psi_c^D(R)$. Therefore, $|\Omega_0| = |R_0|$. Observe that the parity of R is not changed in the revisions. Hence, R_0 is an r -cycle (indeed an r -circuit) in G_{D_0} .

By comparing the two phases of PROJMINMAX we see that its overall time complexity is $O(n^3)$ and its overall space complexity is $O(n^2)$.

Chapter 5

Final Comments

5.A Remarks on Chapter 1

In Chapter 1 we describe a graph model for the study of graphs embedded in surfaces. We show in this thesis that, with this language, certain problems of topological flavor can be formulated in combinatorial terms. The principal advantage of the graph approach is that a 2-dimensional structure is naturally encoded by a colored 1-dimensional structure. An important point in favor of this methodology is that the corresponding concept to topological orientability is the simpler concept of parity; in consequence, orientation-reversing circuit corresponds to odd circuit. The graph definition also provides a geometric-algebraic bridge via Cayley color graphs and Tutte's axioms for a map [25]. Elementary properties of permutations and the definitions introduced yield a simple combinatorial proof that the Euler characteristic of an orientable surface is even; see Theorem 1.7. In Theorem 1.8 we prove a fact about cubic 3-edge-colored graphs which corresponds topologically to the property that no union of a disjoint collection of polygons in G_M whose induced circuits are r -circuits in M^t forms the boundary of a subset of faces in M^t . The model used also leads to the symmetric theory in vertices, faces, and zigzags which motivates the definition of the phial map and of the anti-map, which are maps analogous to the well-known dual map. The dual and

phial maps form the support for the extension of planar duality presented in Chapter 2. To conclude this section, we describe two generalizations of the graph model for a map which merit investigation:

(I) In the definition of a map, we drop the restriction that two of the colors must induce squares. The faithful embedding of a 3-edge-colored cubic graph, with the colors ordered is a natural embedding of a hypergraph: the hyperedges are the polygons induced by the two first colors. They hypervertices are the polygons induced by the two last colors. Each polygon induced by the first and last color is a “face” which, as in the case of graphs, is not an intrinsic concept, because it depends on the particular embedding.

(II) We can try to move one step further in dimension. Consider a 4-regular 4-edge-colored graph, in which the faithful embedding of each component induced by any three of the four colors is in a 2-sphere. We can “fill with 3-space” each one of these 2-spheres, obtaining solid 2-balls. Observe that each bicolored polygon, “filled with 2-space”, occurs twice as 2-faces in the collection of 2-spheres. Thus, we have a natural method for the identification of the boundaries of the 2-balls, thus forming a 3-manifold without boundary. Hence, we can talk about faithful embeddings for the members of the above class of 4-regular, 4-edge-colored graphs, which we call for short *4-graphs*. By, analogy let us call a cubic 3-edge-colored graph a *3-graph*.

The Proof of Theorem 2.0.2 implies that the faithful embedding of a 3-graph is in a 2-sphere iff the cycle space of the 3-graph is generated by its bicolored polygons. An important open question is the following: are there necessary and sufficient graph-theoretical conditions for the faithful embedding of a 4-graph to be in a 3-sphere?

5.B Remarks on Chapter 2

In chapter 2 we introduce rich maps and graphs, as a generalization of planar duality. Loosely speaking, rich graphs are the graphs which have two abstract

duals. Planar graphs are the class of graphs whose members have one abstract dual. Since rich graphs are a natural abstract extension of planar graphs, perhaps some important theorems for the latter extend to the former.

At present, however, more needs to be understood about the nature of rich graphs, perhaps along the topological lines of Theorem 2.9. To us, the more important questions about rich-graphs are the following:

- (i) Are all rich graphs map-rich?
- (ii) is it possible to strengthen Theorem 2.9 so as to provide an algorithmically good characterization of non-map-rich graphs?

To conclude this section we state, without proof, the following graph-theoretic characterization of imbalances in rich maps.

Theorem For a rich map M , a subset $T \subseteq SQ(M)$ is an imbalance in M iff T corresponds to a subset of edges which is a cycle in G_D and the complement of a cycle in G_P . (D and P are the dual and the phial of M , respectively).

5.C Remarks on Chapter 3

In Chapter 3 we solve a generalized version of the Gauss code problem, treating together the cases in which the surface involved is either the plane or the projective plane, under the restriction of 2-face-colorability.

As a generalization we can remove the restriction of 2-face-colorability and try to solve the problem for the projective plane. But as it becomes clear from the text of Chapter 3, this is a completely different problem. More relevant for the present context would be to find an algorithm to determine what is the minimum connectivity of a surface where a realization of a given Gauss code (satisfying the 2-face-colorability restriction) exists. As we showed in Section 3, this would provide an algorithm to determine whether or not a graph with maximum valency 3 is map-rich.

Recently we have asked ourselves how one might algorithmically decide realizability of a Gauss code in the Klein bottle. It seems that the problem is transformed into a linear system of equations in $GF(2)$. However, it looks like the methods which we are using do not extend for the torus. If so, the transition from the Klein bottle to the torus is the breaking point for the techniques that we have available at present.

Another line of generalization of the Gauss code problem is to permit the same symbol to occur an arbitrary number of times in the input sequence. Thus, multiple crossings appear in the curve. This generalization was suggested by Grünbaum [10]. This version of the problem is related to the embedding of hypergraphs.

5.D Remarks on Chapter 4

In Chapter 4 we show that to decide whether a given set of squares of a map is a minimum imbalance is an *NP*-complete problem. We provide a solution for this problem for the class of projective maps. The natural question is: how far can we go? Is it possible to solve the same problem for Kleinian maps? More restrictively, which are the classes of maps, beyond the projective maps, for which the minimax relation described in Theorem 4.5 holds? An apparently unrelated class where it holds is the class of antimaps of planar maps with eulerian graphs. This follows from a result by Seymour [23], strengthening a theorem by Lovasz [16]. This result implies that a minimum transversal of odd coboundaries in bipartite graph has the same cardinality as a maximum disjoint collection of odd coboundaries. For planar graphs the minimal coboundaries correspond to circuits in the planar dual. Thus, Seymour's result implies: for planar eulerian graphs, the minimum transversal of odd circuits is equal in cardinality to a maximum disjoint collection of odd circuits.

To show that the above minimax relation implies (4.5) for the class of antimaps of planar maps with eulerian graphs, observe that it is enough to

establish that for m^t , where $M^\sim$ is an orientable map, the concept of r -circuit is synonymous to odd circuit in $G_M = G_{M^\sim}$. To prove that the latter two concepts are equivalent we show, initially, that for an arbitrary map M , a circuit X in G_M is an r -circuit in M^t iff for some imbalance I (and thus an arbitrary imbalance I), $|X \cap I|$ is odd, where I is considered a subset of eG_M . Let X' be a circuit in C_M such that $\psi_c^M(X') = X$. Let $\{A, B\}$ be a balancing partition for M . Denote by I the imbalance induced by $\{A, B\}$ and by f_I the f_M -edges in the squares of I . Observe that $eG_M \setminus \delta A = f_I$ and that the parity of $|X' \cap f_I|$ is equal to the parity of $|X'|$. Also, $|X' \cap f_I| \equiv |X \cap I| \pmod{2}$. By definition of r -cycle, it follows that X is an r -circuit iff $|X \cap I| \equiv |X'|$ is odd.

Now we conclude that for the antimap of an orientable map, r -circuit is a synonym for odd circuit. If $M^\sim$ is orientable, then ϕ is an imbalance in $M^\sim$ and, by (3.2.1), eG_M is an imbalance in M . Put eG_M in the role of I in the above equivalence. The proof is complete.

Two classes of maps for which 4.5 holds are, therefore, the class of projective maps M with G_M eulerian, and the class of maps M , where G_M is eulerian and $M^\sim$ is planar. These classes are, apparently, unrelated. It would be interesting to find a common generalization of these classes for which (4.5) holds.

Bibliography

- [1] Bondy, A. and Murty, U., *Graph Theory with Applications*, American Elsevier (1976).
- [2] Cook, S.A., *The Complexity of Theorem Proving Procedures*, Proc. 3rd Annual ACM Symposium on Theory of Computing, 151–158.
- [3] Coxeter, H.S.M., *Regular Polytopes*, Dover (1973).
- [4] Edmonds, Jack, *A combinatorial Representation for Polyhedral Surfaces*, Notices of Amer. Math. Soc. 7 (1960) 646.
- [5] Edmonds, Jack, *On the surface Duality of Linear Graphs*, Journal of R.N.B.S. vol. 69B (1965) 121–123.
- [6] Garey, M.R. and Johnson, D.S., *Some Simplified NP-Complete Problems*, Proc. 6th Annual ACM Symposium on Theory of Computing, 47–63.
- [7] Giblin, P.J., *Graphs, Surfaces and Homology*, Chapman and Hall (1977).
- [8] Godement, R., *Algebra*, Hermann, Paris (1968).
- [9] Gross, I. and Magnus, W., *Groups and Their Graphs*, New Math. Library, 14, Math Assoc. of America (1964).
- [10] Grünbaum, B., *Arrangement and Spreads*, Conf. Board of the math. Sciences Regional Conf. Series in Math. no. 10.

- [11] Henle, M., *A Combinatorial Introduction to topology*, Victor Klee, Freeman and Company (1979).
- [12] Hoffman, Peter, *Any Embedding is Combinatorial*, Mimeographed Notes.
- [13] Karp, Richard M., *Reducibility Among Combinatorial Problems*, Miller and Thatcher ed (1972) 85–104.
- [14] Lins, Sóstenes, *A Sequence Representation for Maps*, To appear in Discrete Mathematics.
- [15] Lins, Sóstenes, *A Minimax Theorem on Circuits in Projective Graphs*, To appear in the JCT.
- [16] Lovasz, L., *2-Matchings and 2-Covers of Hypergraphs*, Acta Math. Acad. Scientiarum Hungaricae, Tomus 26 (3–4), (1975), 433–444.
- [17] Lovasz, L., Marx, M., *A Forbidden Substructure Characterization of Gauss Codes*, Acta Sci. Math. 38 (1976) 115–119.
- [18] Ringel, G., *Map Color Theorem*, Springer-Verlag (1974).
- [19] Rosenstiehl, P., *Solution Algebrique du Probleme de Gauss sur la Permutation des Points d'Intersection d'une ou Plusieurs Courbes Fermées du Plan*, C.R. Acad. Sc. Paris, t.283 (11 octobre 1976) 551–553.
- [20] Rosenstiehl, P., *Characterization des Graphes Planaires par une Diagonale Algebrique*, C.R. Acad. Sc. Paris, t.283 (4 octobre 1976).
- [21] Read, R.C. and Rosenstiehl, P., *On the principal Tripartition of Graphs and Binary Matroids*, annals of Discrete Math. vol. 3 (1978).
- [22] Schubert, H., *Topology*, Macdonald Technical and Scientific, 1968 (Translation from the German).
- [23] Seymour, P.D., *On Odd Cuts and Plane Multicommodity Flows*, To appear in JCT (series B).

- [24] Shank, H., *The Theory of Left-Right Paths*, Combinatorial Mathematics III Lecture Notes no 452, Springer-Verlag (1975).
- [25] Tutte, W.T., *What is a Map?* New direction in the theory of graphs, F. Harary ed. (1971).
- [26] White, Arthur T., *Graphs, Groups and Surfaces*, North-Holland/American Elsevier (1973).
- [27] Whitney, H., *Non-Separable and Planar Graphs*, Trans. Amer. Math. Soc. 34 (1932) 339–362.
- [28] Younger, D.H., *Embeddability of Graphs in the Projective Plane*, Personal Communication.
- [29] Youngs, J.W.T., *The Solution of the Heawood Conjecture*, Graph Theory and Theoretical Physics, Frank Harary Ed. 313–354, Acad. Press (1967).
- [30] Walsh, T., Mimeographed notes on Map Theory for lectures given at the University of Waterloo during May 1979.